

RESEARCH ARTICLE

Comparative gene expression profiles for highly similar aggressive phenotypes in male and female cichlid fishes (*Julidochromis*)

Molly Schumer, Kavita Krishnakant and Suzy C. P. Renn*

Department of Biology, Reed College, 3203 S. E. Woodstock Boulevard, Portland, OR 97202, USA

*Author for correspondence (renns@reed.edu)

Accepted 5 July 2011

SUMMARY

Julidochromis marlieri and *Julidochromis transcriptus* are two closely related Tanganyikan cichlids that have evolved different behavior and mating strategies since they diverged from their common ancestor. While *J. transcriptus* follows the ancestral pattern of male dominance, male-biased sexual size dimorphism and territoriality, the pattern is reversed in *J. marlieri*. In *J. marlieri*, females show all of these behavioral and morphological characteristics. This raises the question of whether female *J. marlieri* achieve the dominant phenotype by expressing the same genes as *J. transcriptus* males or whether novel brain gene expression patterns have evolved to produce a similar behavioral phenotype in the females of *J. marlieri*. This study used cDNA microarrays to investigate whether female *J. marlieri* and male *J. transcriptus* show conserved or divergent patterns of brain gene expression. Analysis of microarray data in both species showed certain gene expression patterns associated with sex role independent of gonadal sex and, to a lesser extent, gene expression patterns associated with sex independent of sex role. In general, these data suggest that while there has been substantial divergence in gene expression patterns between *J. transcriptus* and *J. marlieri*, we can detect a highly significant overlap for a core set of genes related to aggression in both species. These results suggest that the proximate mechanisms regulating aggressive behavior in *J. transcriptus* and *J. marlieri* may be shared.

Supplementary material available online at <http://jeb.biologists.org/cgi/content/full/214/19/3269/DC1>

Key words: sex-specific behavior, brain gene expression, isotocin, AVT.

INTRODUCTION

Parental care demands an investment of parental resources, and represents a trade-off in fitness for each parent (Trivers, 1972). In biparental species, males and females share the fitness burden of parental care but, despite this, males and females frequently differ in the amount and nature of parental care they provide, showing strong biases such that sex-specific behaviors develop. For a wide range of biparental species that exhibit sex-specific behaviors, it is common to find males engaged in territorial defense while females are involved in more direct care of offspring [e.g. birds (Burger, 1981; Creelman and Storey, 1991; Fraser et al., 2002); and fish (Keenleyside and Bietz, 1981; Lavery and Reeb, 1994; Wisenden, et al., 1995; Itzkowitz et al., 2001; Itzkowitz et al., 2003; Itzkowitz et al., 2005)].

In a minority of species, the conventional bias of sex-specific behavior is reversed such that females engage in aggressive behavior while males perform the majority of direct parental care [e.g. pipefish (Berglund and Rosenqvist, 2003); and Wattled Jacanas (Emlen and Wrege, 2004)]. Reversal of sex-specific behavior in these species is often assumed to be the result of females having a higher potential reproductive rate (Eens and Pixten, 2000; Cluttonbrock and Vincent, 1991), ultimately resulting in stronger sexual selection on females. This ultimate explanation has rarely been evaluated at a genetic level (Jones et al., 2001) but is well supported by studies that show plasticity of sex-specific behavior under ecological fluctuations that reverse the direction of sexual selection (Forsgren et al., 2004; Gwynne, 1985). Even when these ecologically based proximate causes of reversed sex-specific behavior are identified, the

mechanistic explanations often remain elusive. Despite investigation of hormone levels [e.g. pipefish (Mayer et al., 1993); and birds (reviewed in Eens and Pixten, 2000)] and gene expression (Voigt and Goymann, 2008), it has been difficult to determine the extent to which the mechanisms that underlie aggressive behavior in females are similar to those that underlie aggressive behavior in males. Species that show reversed sex-specific parental care behavior provide the opportunity to examine the proximate mechanisms that underlie reversed phenotypes.

The African cichlids provide an excellent model with which to address the proximate mechanistic explanations of sex-specific behavior. Their adaptive radiation has been accompanied by a vast amount of behavioral diversification, including several independent transitions in parental care strategy (Goodwin et al., 1998). A number of studies demonstrate that within the family Cichlidae biparental behaviors are divided into stable sex-specific behavioral phenotypes. The existing body of research suggests that biparental cichlids generally show sex-biased behavior, with females performing the majority of parental care and males utilizing their increased size for territory and brood defense (Keenleyside and Bietz, 1981; Itzkowitz, 1984; Kuwamura et al., 1986; Kuwamura et al., 1997). The cichlid genus *Julidochromis* is composed of biparental substrate brooders (Konings, 1998) that show variation in specialized parental care such that several species in this genus exhibit the conventional pattern of sex-specific behavior while *Julidochromis marlieri* shows reversed sex-specific behavior (Barlow and Lee, 2005). *Julidochromis marlieri* females are large and highly aggressive (Yamagashi and Kohda, 1996; Barlow and Lee, 2005) (K. Wood

and S.C.P.R., unpublished), and *J. marlieri* males spend significantly more time in the nest when eggs are present (Schumer, 2009). Although estimates vary, according to recent phylogenetic analysis, *Julidochromis transcriptus* and *J. marlieri* diverged ~3.7 million years ago (Sturmbauer et al., 2010). In our present study, the molecular mechanisms that underlie sex-specific behavior in *J. marlieri* males and females were directly compared with those of another *Julidochromis* species, *J. transcriptus*, which has conventional sex-specific behavior (Awata et al., 2006; Schumer, 2009). The present study focuses on investigating whether the proximate mechanisms underlying dominant and subordinate behavior are the same in *J. transcriptus* and *J. marlieri* or whether these mechanisms differ due to reversal of sex-specific behavior in *J. marlieri*.

Recent advances in the ability to identify the genes involved in complex phenotypes means that we are no longer constrained to mutant screens in model organisms [circadian rhythm in *Drosophila* (Konopka and Benzer, 1971); chemotaxis in *Caenorhabditis elegans* (Dusenbery, 1980); and mating in *Drosophila* (Yamamoto et al., 1997)] and can even move beyond the identification of natural allelic variation [foraging behavior in *Drosophila* (Osborne et al., 1997)] to take on a genome-wide comparative approach. The development of reliable methods to globally assay gene expression levels and associate these gene expression patterns with behavior has revolutionized our understanding of the relationship between gene expression and complex behaviors such as aggression (Edwards et al., 2006), addiction (Lewohl et al., 2000; Rhodes and Crabbe, 2005; Buitenhuis et al., 2009), maternal care (Gammie et al., 2007), social behavior (Whitfield et al., 2003) (reviewed by Robinson et al., 2008), and even mate choice (Cummings et al., 2008). Among African cichlids, microarray techniques have been used to examine the basis of a variety of behaviors including dominance and subordinate behavior in male *Astatotilapia burtoni* (Renn et al., 2008), various phenotypes in *Neolamprologus pulcher* breeding groups (Aubin-Horth et al., 2007) and different mating strategies in species from the Ectodini clade (Machado et al., 2009). Combined, those studies have identified a number of genes whose expression strongly correlates with aggressive behavior or with sex, thus demonstrating that certain sex-specific behaviors in cichlids are accompanied by differences in gene expression.

Few studies to date have compared gene expression patterns between similar phenotypes in different species. Microarrays can be used to ask these comparative questions but become less useful for comparing phenotypes between species as genetic divergence increases (Machado et al., 2009; Renn et al., 2004; Cohen et al., 2007). In closely related species, microarrays have been successfully used to investigate whether differences in phenotype are reflected by changes in phenotype-specific gene expression. Honeybee species have slightly different species-specific foraging dances; gene expression profiles of three species producing the foraging dance revealed striking similarities between species but also identified a number of species-specific differences in gene expression in genes related to motor control and synaptic activity (Sen Sarma et al., 2009). Comparative work investigating the genes underlying a similar foraging behavior in honeybees and paper wasps showed that a small set of genes was shared between these independently evolved phenotypes (Toth et al., 2010). Such studies expand the potential questions posed in microarray studies from those focused on assessing gene expression differences within species to broader questions about the evolution of phenotypic differences between species.

The main aim of the present study was to identify the proximate mechanisms for the variation in sex-specific roles observed among species of the genus *Julidochromis*. We asked whether when a female *J. marlieri* achieves the conventionally male-like dominant phenotype, she does so through a pattern of neural gene expression similar to that of a *J. transcriptus* male or whether this reversal of the behavioral phenotype is orchestrated through a novel pattern of brain gene expression. The former would suggest the co-option of homologous molecular mechanisms to produce a male-like dominant phenotype in these females whereas the latter would suggest convergent evolution to produce superficially similar behaviors in the opposite sexes of these two lineages.

Using cDNA microarrays we characterized differences in brain gene expression between sex within each species to identify genes that are associated with sex-specific behavior in each species. We then compared the list of female-biased genes in *J. marlieri* with the list of male-biased genes in *J. transcriptus* to ask whether female aggression in *J. marlieri* relies upon the same genetic mechanisms as male aggression in *J. transcriptus*. We also compared the list of male-biased genes from *J. marlieri* with the list of female-biased genes from *J. transcriptus* to identify genes that are associated with subordinate behaviors, regardless of sex and species. Keeping in mind that downregulation of gene expression level may be equally important for the observed phenotypes, the observed regulation of these subordinate-biased genes may also be important for the dominant phenotype. Although the majority of gene expression was both phenotype and species specific (~10% of differentially expressed genes shared between phenotypes), we found convincing evidence for a core set of genes regulating aggressive behavior with conserved expression between species.

MATERIALS AND METHODS

Animal husbandry

Fish were maintained at pH 8.0 and 650 μ S salinity with a 5% daily water change, and light cycle including 11.5 h of light with a 30 min fade for dawn and dusk to mimic conditions in nature. Pairs of *Julidochromis marlieri*, Poll 1956, and *Julidochromis transcriptus*, Matthes 1959, were allowed to form naturally from mixed male–female populations in 113.6 l tanks. Two individuals were considered to have formed a ‘pair’ when they were observed over multiple days to have established a territory together. After a pair was established it was transferred to a 113.6 l observation tank that consisted of a neighboring pair of the same species separated by a transparent acrylic divider. Each pair was provided with ~2 cm of gravel substrate for digging and an artificial nest crevice. The nests were made of two clay tiles (15 \times 15 \times 1 cm) with an entrance width of 8 cm, designed to mimic the caves and crevices that are used as nests in the wild (Awata et al., 2006). *Julidochromis transcriptus* and *J. marlieri* stocks had been maintained in-house for three years at the time of the study. First filial generation *J. transcriptus* individuals were obtained from Atlantis Tropical Fish Hatchery (Gardiner, NY, USA), and *J. marlieri* individuals were obtained from George Barlow, Berkeley (CA, USA), formerly at UC Berkeley (age uncertain).

All experiments were carried out according to animal protocol IACUC no. 1032007.

Behavioral observations

Eight pairs (four pairs of each species) were checked every morning between 08:00 h and 10:00 h for eggs and observed weekly using the event recorder Jwatcher (Blumstein and Daniel, 2007). Ten-minute behavioral observations were used to confirm that individuals

were exhibiting the sex-specific behavior typical of their species. Males and females within a pair were observed simultaneously. The ethogram included event measures of aggressive behaviors (attack intruder, attack mate, approach intruder) and parental care behaviors (egg cleaning) as well as duration measures of parental care behavior (time spent in the nest). The total number of observations for each pair varied based on when eggs were laid but ranged from 3 to 11 observations. Due to the differences in the number of observations, a generalized linear model was used with the lme4 package in R (<http://cran.r-project.org/web/packages/lme4/index.html>) to test for differences in male and female behavior in each species and for overall differences in behavior between species. Two post-spawning behavioral observations were conducted for each pair, one on the day of egg laying and one 24 h later.

Sample collection

Twenty-four hours after a pair had laid eggs, the male and female were simultaneously removed from the observation tank, anesthetized using MS-222 (160 mg of MS-222 in 500 ml cichlid salt water, pH 8.0) and killed according to protocol (IACUC #2007.103). This time point was chosen in order to ensure synchrony of reproductive cycles and capture parental care. After the fish had been anesthetized, two researchers simultaneously dissected the two fish. Following decapitation, the brain was immediately removed and stored in 1 ml RNAlater (Ambion, Austin, TX, USA). Brains were stored at 4°C for 24 h and then at -20°C. Gonads were removed and stored in 0.5 ml paraformaldehyde at 4°C. Samples were collected from four pairs of each species.

RNA extraction and microarray hybridization

Brains were homogenized (Tissue tearor, Biospec Products, Bartlesville, OK, USA), and total brain RNA was isolated using a standard Trizol (Invitrogen, Carlsbad, CA, USA) protocol adapted for use with phase-lock gel tubes (Eppendorf, Westbury, NY, USA). RNA was quantified with a Nanodrop 1000 (Thermo Scientific, Wilmington, DE, USA). Each total RNA sample, ~1000 ng, was linearly amplified with a single round of Amino Allyl MessageAmp II aRNA amplification (Ambion), according to manufacturer's protocol (yield 31- to 74-fold). The antisense amplified RNA samples, 4 µg each, were then dye-coupled with Cy3 and with Cy5 and quantified with the Nanodrop 3300 fluorospectrometer (Thermo Scientific) prior to storage at -80°C. One limitation of using whole brain RNA is the possibility that this method will fail to detect certain changes in gene expression, due to discrete anatomical localization or counter regulation in differing brain regions; therefore, subtle regulation of gene expression may be missed by our technique (Filby et al., 2010).

In each hybridization, equal quantities of Cy3 and Cy5 samples (1.5–2.5 µg) were combined in hybridization buffer (Ambion SlideHyb buffer #1) and applied to the second-generation cDNA array (GEO platform ID: GLP6416) designed for the African cichlid *A. burtoni* (Renn et al., 2004; Salzburger et al., 2008), which contains ~20,000 features, representing ~16,000 unique sequences (including ~35% anonymous features) derived from brain, skin and mixed organ libraries from individuals of all ages, sexes and phenotypes, and has been validated for a variety of cichlid species (Renn et al., 2004; Aubin-Horth et al., 2007; Machado et al., 2009). Cross-species comparative genomic DNA hybridization for these *Julidochromis* species identified fewer than 10 microarray features that are substantially influenced by sequence or copy number differences between these species (data not shown). Given the within-species design, even these features should not affect the interpretation of




Fig. 1. Experimental hybridization loop design used for *Julidochromis transcriptus* (A) and *Julidochromis marlieri* (B). Each individual was hybridized to two or three conspecifics of the opposite sex. The head of the arrow represents a sample labeled in Cy5 and the tail of the arrow represents a sample labeled in Cy3.

results. For each species, the eight individuals were compared between sex and within species (Fig. 1) in a modified loop design (Churchill, 2002) incorporating dye swaps for greater statistical power. Each sample was used for 2–3 hybridizations, one of which was to their mate.

Microarrays were incubated at 48°C for 16–18 h and then rinsed through three successive wash buffers and spun dry prior to scanning on a GenePix 4000B scanner (Molecular Devices, Sunnyvale, CA, USA) with GenePix 5.1 imaging program. Poor spot morphology and technical artifacts were flagged manually. These raw data were imported into R software (v.2.12) (R Development Core Team, 2006) for analysis with the Linear Models for Microarray Data package (LIMMA) (Smyth, 2004). Array features were filtered for poor spot morphology, hybridization artifacts and low intensity (<2 s.d. above local background). Background subtracted intensities were normalized within array using the print tip LOESS method, and replicate features representing the same DNA sequence were averaged and weighted. Intensities were grouped for analysis using the lmFit function adapted to calculate exact *P*-values in the presence of missing data (A. Jones and S.C.P.R., unpublished), and evaluated for sex-specific bias using the empirical Bayes method (eBayes) (Smyth, 2004). Only genes surviving quality filters in sufficient replicates for analysis in both species were considered for the analysis of differential regulation. Genes deemed significant at $P < 0.05$ were investigated using a gene index annotation developed for the *A. burtoni* cDNA array (Renn et al., 2004) (http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=a_burtoni); no multiple hypothesis testing correction was used due to small sample size. Lists of significantly regulated genes for each phenotype were compared using one-line perl scripts provided by FAS Center for Systems Biology (<http://sysbio.harvard.edu/csb/resources/computational/scriptome/>). All between-species comparisons were done through gene lists rather than statistical comparison, due to the fact that no between-species hybridizations were performed in order to prevent hybridization biases caused by species-specific sequence differences. To test for significant overlap between different gene lists, a Fisher's exact test was used. All raw and processed data are available at the GEO database [(www.ncbi.nlm.nih.gov/projects/geo/) Series: GSE23094 arrays, GSM569290–GSM569309].

Real-time quantitative PCR (RT-qPCR) methods

Gene-specific primers were designed for four genes (Table 1) using sequence information from *A. burtoni*: parvalbumin (DY626384),

Table 1. Primer sequences for the candidate genes and housekeeping gene verified by quantitative PCR

Gene name	Forward primer (5'→3')	Reverse primer (5'→3')
Isotocin	GGTCTCACTGCAAGCTCTCG	CAGGGGAAAAGAGCAAAACC
Parvalbumin	GCACATTCGACCACAAGAAG	ATGAAACCGCTCTTTGCTCTG
MAP-1B	CTGTTGCCACACAAAAGGAC	TGGTCTTCGGAGAGGATTTG
GAPDH	CCTCCATCTTTGATGCTGGT	GTGCATGTACAGCAGCAGGT

MAP-1B, microtubule-associated protein-1B; GAPDH, glyceraldehyde 3-phosphate dehydrogenase.

isotocin precursor (DY631081), microtubule-associated protein-1B (MAP-1B, DY628922) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH). Primers were designed using Primer3Plus and optimized using IDT Oligo Analyzer (<http://www.idtdna.com>). Primer specificity (single band of predicted size) was verified for both *Julidochromis* species using genomic DNA (gDNA) as a template. RNA samples were available in sufficient quantity for technical validation by RT-qPCR using all eight *J. transcriptus* individuals and seven of the eight *J. marlieri* individuals used in the original hybridizations. DNA template was synthesized using oligo (dT) for total RNA and random hexamers for amplified antisense RNA (iScript, Biorad, Hercules, CA, USA). Per reaction volumes were: 2 µl 5× iScript reaction mix, 1 µl primer, 0.5 µl iScript reverse transcriptase, 25 ng µl⁻¹ of RNA from each individual, and water (added to bring the final reaction volume to 10 µl). For an oligo (dT) primed reaction, the mix was incubated at 42°C for 90 min and then 85°C for 5 min. For a random hexamer primed reaction, the mix was incubated at 25°C for 5 min, followed by an incubation at 42°C for 30 min and finally at 85°C for 5 min. Reaction conditions for RT-qPCR were optimized by creating a standard curve using different primer and template concentrations; efficiency for all primers ranged from 90% to 100%. RT-qPCR (Opticon; Biorad) was performed on cDNA using GAPDH for normalization. Per reaction volumes were: 5 µl 2× ImmoMix (Bioline, Taunton, MA, USA), 0.15 µl 50× SYBR Green, 5 µmol l⁻¹ of forward primer and reverse primer (Integrated DNA Technologies, San Diego, CA, USA), 2.85 µl nanopure water, and 1 µl template cDNA. All qPCR reactions were performed as follows: 1 cycle of 95°C for 10 min, 35 cycles of 94°C for 10 s, 57–60°C for 20 s, 72°C for 15 s, followed by a hold at 72°C for 2 min. At the end of the reaction a melt curve analysis was performed for each sample. Each sample was run in triplicate and each plate contained no RT controls. A linear model was generated using the `lm` function in R to look for significant correlation in gene expression between qPCR and microarray results.

RESULTS AND DISCUSSION

Behavioral observations confirmed that *J. transcriptus* pairs show conventional sex-biased behavior whereas in *J. marlieri* this behavior is reversed. This allowed us to assess our central question: is aggressive behavior regulated by similar or divergent gene expression mechanisms in both *Julidochromis* species despite reversal of sex-specific behavior? Overall, microarray results support the conclusion that gene expression in *J. marlieri* has diverged from *J. transcriptus*. However, we also found a core set of genes associated with the aggressive phenotype in both species, some of which are confirmed by qPCR. We found evidence of sex-biased gene expression and phenotype-specific gene expression that included a number of candidate genes previously implicated in aggression or parental care in a variety of species (Table 2). The territorial/aggressive phenotypes of both species showed the greatest number of differentially expressed genes. These dominance-biased

genes tended to be those of greater fold change, and overlap between the gene lists for the two aggressive phenotypes of each species was highly significant. Functional inference can be drawn from the available annotations (Table 2) (http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=a_burtoni), and regulated genes with no known homolog represent novel candidates for the molecular control of behavior.

Sex-biased differences in behavior

The pairs that bonded conformed to expected sexual size dimorphism for both species. *Julidochromis transcriptus* males weighed more (2× mean; $P=0.006$) and were larger (1.3× mean; $P=0.008$) than their mates whereas *J. marlieri* females weighed more (2.6× mean; $P<0.001$) and were larger (1.5× mean; $P<0.001$) than their mates. There was no significant difference in gonadosomatic index (GSI) between species.

In order to confirm species-specific patterns of sex-biased behavior, approach conspecific, attack conspecific and attack mate behaviors were totaled to quantify aggression. Due to the different numbers of observations per pair, a generalized linear model was used for each species to assess whether variation in behavior was best explained by sex, mate or individual. Based on this analysis, *J. transcriptus* males were found to be significantly more aggressive than females ($P<0.0001$, Fig. 2). Conversely, *J. marlieri* females were significantly more aggressive than males ($P=0.00083$, Fig. 2).

We examined the post-spawning time point based on the assumption that sex-specific behavior would be more pronounced after eggs were laid, because maximizing effective parental care is one of the hypothesized functions of sex-specific behavior. As predicted, more parental care was provided by *J. transcriptus* females ($P=0.057$) and by *J. marlieri* males ($P=0.042$) as measured by the time in the nest after spawning. This suggests that the subordinate individual of each species provides the majority of parental care. The measure of direct parental care, egg cleaning, which occurred infrequently, did not show a statistically significant sex bias as had been predicted. While *J. marlieri* post-spawning females tended to increase aggression (Fig. 2), as predicted, post-spawning *J. transcriptus* males tended to reduce aggression. Therefore, the sampling time point for gene expression may not have measured the peak in aggressive behavior in *J. transcriptus*. However, as post-spawning *J. transcriptus* males stayed closer to the nest, there was reduced opportunity to capture aggressive behavior with our ethogram. Although untested, we predict that behavioral assays including intruder introductions and territory challenges might reveal an increase in post-spawning aggression for *J. transcriptus* males.

Based on a generalized linear model including both species, *J. marlieri* females were significantly more aggressive than *J. transcriptus* males in both control ($P=0.0484$) and post-spawning conditions ($P=0.0046$) while *J. transcriptus* individuals spent more time in the nest with eggs compared with *J. marlieri* individuals ($P=0.0669$). Overall, the behavioral results confirmed that *J. marlieri*

Table 2. Candidate genes for controlling behavior upregulated in *Julidochromis transcriptus* and *Julidochromis marlieri* males and females

Group	Annotation	Potential function
<i>J. transcriptus</i> females	Homolog of GHRH precursor	GHRH is expressed in the preoptic nucleus ^a
	Early growth response 1	Protein expressed throughout the brain, associated with stimulus response ^b
<i>J. transcriptus</i> males	Isotocin	AVT paralog ^c
	Gonadotropin α-subunit	Aggression, gonad development ^d
	Parvalbumin-1 precursor	Protein associated with GABA neurons ^e
	Somatotropin	Expressed in dominant <i>Astatotilapia burtoni</i> males ^f
	Prolactin precursor-1	Implicated in parental care behavior ^g
<i>J. marlieri</i> females	Isotocin	AVT paralog ^c
	Parvalbumin-1 precursor	Protein associated with GABA neurons ^e
	Arginine vasotocin preprohormone	Associated with aggressive behavior ^h
	Aromatase homolog	Aromatase is associated with aggressive behavior ⁱ
<i>J. marlieri</i> males	Prolactin precursor-1	Implicated in parental care behavior ^g
	Gonadotropin α-subunit	Aggression, gonad development ^d
	Somatolactin	Expressed in dominant <i>A. burtoni</i> males ^f

Genes that were also significantly upregulated for another phenotype are indicated in bold.

GHRH, growth hormone-releasing hormone; AVT, arginine vasotocin; GABA, γ -aminobutyric acid.

^aMarivoet et al., 1988; ^bBurmeister and Fernald, 2005; ^cReviewed in Insel and Young, 2000; ^dPrice et al., 2003; Millesi et al., 2002; Foran and Bass, 1999;

^eCelio, 1986; ^fRenn et al., 2008; ^gReviewed in Schradin and Anzenberger, 1999; ^hRenn et al., 2008; Aubin-Horth et al., 2007; Santangelo and Bass,

2006; ⁱRenn et al., 2008; S.C.P.R. and H. A. Hoffman, unpublished; Soma et al., 1996; ^jRenn et al., 2008.

females and *J. transcriptus* males are more aggressive while *J. marlieri* males and *J. transcriptus* females spend more time in direct contact with eggs.

Sex-biased gene expression

Males and females of the same species share the majority of their genome but can differ substantially in their sex-specific behavior, suggesting an important role of gene expression. In some studies, sex-specific gene expression represents the majority of within-species variation in gene expression (Yang et al., 2006; Jin et al., 2001) while other studies find few sex-specific differences in gene expression, even when adults are sexually and behaviorally dimorphic (Santos et al., 2008). Similar to the latter example, the present study found that despite the sexual dimorphism in behavior observed in both *Julidochromis* species, males and females differed in expression for a minority of the genes analyzed by microarray (~7.5% in *J. transcriptus*, ~9.3% in *J. marlieri*). Although few in number, in part owing to small sample size, the genes that were differentially regulated between males and females in each species present good candidates for the molecular basis of behavioral and physiological differences between males and females in *J. transcriptus* and *J. marlieri*.

We found a number of genes that were differentially regulated by sex in each species ($P < 0.05$). Out of 9528 genes analyzed in both species (5111 with sequence information), 301 showed female-biased expression and 412 genes showed male-biased expression in *J. transcriptus*. In *J. marlieri*, 529 genes showed female-biased expression and 356 genes showed male-biased expression. Approximately 15% of the genes analyzed were differentially regulated by one of the four phenotypes. The species difference in the number of regulated genes is not likely due to statistical power as witnessed by the similarity in gene expression level 50 (GEL50) values (Clark and Townsend, 2007) for these two experiments (*J. transcriptus* 0.2397; *J. marlieri* 0.2129). Gene lists from males and females of both species were compared to identify any genes that were sex specific in both species. Genes within these sex-specific modules may be related to reproductive physiology. There was a

trend towards more genes in the female module (24 genes) than expected by chance but not in the male module (21 genes) (Fisher's exact test: female, $P = 0.07198$; male, $P = 0.1426$, Fig. 3). The 45 sex-biased genes consistent between the two species are much fewer than the genes that were found to be phenotype specific (see below). Similarly, a recent study of monogamous and polygynous cichlids found that most differences in gene expression were both sex and species specific, with few genes showing sex-specific gene expression across species (Machado et al., 2009). In a previous review on the topic, Ellegren and Parsch (Ellegren and Parsch, 2007) reported that many of the genes that show sex-specific expression within species are those that are most sequence diverged between species, suggesting functional and therefore likely expression divergence. Therefore, based on previous research, it is not surprising that so few genes were sex specific across species.

A number of the sex-biased genes have a known role in reproduction and mating behavior. Gonadotropin-releasing hormone (GnRH), which is involved in sexual differentiation and mating behavior in organisms from fish to mammals, was strongly male biased in both *Julidochromis* species. While GnRH has been implicated in dominance because it is upregulated in *A. burtoni* territorial males compared with non-territorial males (Renn et al., 2008), elevated GnRH in territorial male *A. burtoni* may be related to gonad growth (see Soma et al., 1996) rather than behavior *per se*. In order to fully understand the role of GnRH in *Julidochromis*, it would be necessary to assess sex-specific expression throughout the reproductive cycle. In many species, including *A. burtoni* females, GnRH varies across reproductive state (White and Fernald, 1993); nonetheless at the time point assayed here, it is interesting to note that GnRH is expressed at higher levels in males. Prolactin, another male-biased gene in both *Julidochromis* species, has been linked to parental care behavior in a range of species [e.g. birds (Garcia et al., 1996); and rats (Bridges et al., 1990)]. This male bias is at first surprising given that females show greater parental care than males in *J. transcriptus*. However, despite the fact that subordinate individuals in *Julidochromis* spent the most time with the eggs, it is known that males and females




Fig. 2. Mean number of aggressive behaviors performed by individuals in a 10 min period during pre- (>3 observations) and post- (2 observations) mating conditions. Male (black) and female (gray) are stacked within pair with symbol size indicating the relative size. *Julidochromis transcriptus* males had significantly higher levels of aggressive behavior before spawning ($P < 0.0001$) but not after spawning ($P = 0.448$) whereas *J. marlieri* females were significantly more aggressive both before ($P = 0.00083$) and after ($P < 0.0001$) spawning.

of both species provide some parental care. Therefore, the male bias in expression may be consistent with the observation that in fish prolactin is particularly important for paternal rather than maternal care (Schradin and Anzenberger, 1999). The bias in prolactin expression in our present study of monogamous species is inconsistent with findings from a related cooperative breeding cichlid species (Bender et al., 2008).

Phenotype-specific gene expression

The present study is among the first to consider whether similar phenotypes in related species are the product of similar gene expression profiles or whether these phenotypes are regulated by different mechanisms. Specifically, we asked whether aggressive behavior is regulated by similar or divergent gene expression mechanisms in both *Julidochromis* species despite reversal of sex-specific behavior. Other studies addressing similar questions have reached varying conclusions. While two limnetic salmonid species show similar activation of genes important for survival in that niche (Derome and Bernatchez, 2006), few genes show consistent activation according to mating system among another cichlid clade (S.C.P.R. and H. A. Hofmann, unpublished).

In order to address the question of phenotype-biased gene expression, we compared gene lists instead of using direct hybridization to reduce problems associated with interspecific hybridization on the microarray. Dominance-related genes were identified as those that were highly expressed in the aggressive male phenotype in the conventionally sex-biased *J. transcriptus* and that were also highly expressed in the aggressive female phenotype in the reversed sex-biased species *J. marlieri* (Fig. 3). Aggressive *J. transcriptus* males and *J. marlieri* females shared increased expression of 76 dominance-related genes (34 with sequence information), significantly more than expected by chance (Fisher's exact test: $P < 2.2e^{-16}$). Conversely, 24 subordinate-related genes were identified as the intersection of the gene lists for highly expressed genes in *J. marlieri* males and in *J. transcriptus* females (16 with




Fig. 3. Venn diagram representing sex-biased gene expression ($P < 0.05$) for *Julidochromis transcriptus* (dark blue, male; pink, female) and *Julidochromis marlieri* (light blue, male; red, female). Overlapping regions show shared sex-specific and phenotype-specific biases.

sequence information), also significantly more than expected by chance ($P = 0.00049$). The fact that only a proportion of the genes that are associated with either social phenotype in each species are shared between species (Fig. 3) suggests that, since *J. marlieri* and *J. transcriptus* diverged from their common ancestor, other changes have also evolved in the gene expression patterns regulating these behaviors. Although few in number, it is notable that both of the phenotype-specific modules had highly significant overlap whereas the sex-specific modules did not. This leads to the exciting interpretation that the few genes that do show phenotype-specific gene expression across species are in fact major regulators of behavior. If the genes that are associated with dominant and subordinate phenotypes in both species are major regulators of behavior, this would support the hypothesis that the sex-specific phenotypes in *J. transcriptus* and *J. marlieri* are controlled by conserved mechanisms. The dominance module (i.e. the overlapping list of genes that are both upregulated in *J. marlieri* females and in *J. transcriptus* males) contained more genes than the opposing subordinate related module or sex-related module, and the overlap between gene lists for the dominance module was more significant than that for any other module ($P < 2.2e^{-16}$). Furthermore, an examination of gene expression patterns (Fig. 4) reveals that many of the most significantly regulated genes in *J. transcriptus* males and *J. marlieri* females are part of the dominance module. These patterns of shared regulation in *J. transcriptus* and *J. marlieri* suggest that the functional mechanisms regulating dominance are conserved between species.

Identified dominance-related genes

Following from the overarching hypothesis regarding gene expression patterns, the individual genes within the dominance- and subordinate-related modules are of great interest as potential key regulators of behavior. Two such genes, highly expressed in both male *J. transcriptus* and female *J. marlieri*, are already known to be involved in brain function. Isotocin, a paralog of arginine vasotocin (AVT) and a homolog to mammalian oxytocin, is a candidate for the regulation of dominant and aggressive behavior in *Julidochromis*. Although the few studies that have investigated a role for isotocin in aggressive behavior have found no correlation (Santangelo and Bass, 2006), its expression pattern here suggests otherwise. Parvalbumin, another identified gene in the dominance




Fig. 4. Volcano plots of sex-specific expression. All colored spots represent significant regulated genes ($P < 0.05$). (A) *Julidochromis marlieri* expression level with significantly biased gene expression indicated for *J. marlieri* (colors as in Fig. 3). (B) *Julidochromis transcriptus* expression level with *J. marlieri* significantly biased gene expression highlighted. (C) *Julidochromis transcriptus* expression level with *J. transcriptus* significantly biased gene expression indicated (colors as in Fig. 3). (D) *Julidochromis marlieri* expression level with *J. transcriptus* significantly biased gene expression highlighted.

module, is expressed in GABAergic neurons (de Almeida et al., 2005), which have been linked to aggressive behavior in a number of mammalian species. The mechanism by which parvalbumin expression might alter γ -aminobutyric acid (GABA) activity to influence aggression is not clear.

AVT was a gene predicted to be upregulated in both *J. transcriptus* males and *J. marlieri* females based on extensive evidence supporting a role for AVT in aggressive behavior. However, AVT was significantly upregulated only in *J. marlieri* females ($P = 0.0164$) but not in *J. transcriptus* males ($P = 0.349$). Although there are dominant phenotypes common to the two *Julidochromis* species, there are also species-specific differences in aggressive behavior; in both pre- and post-spawning conditions *J. marlieri* females were significantly more aggressive than *J. transcriptus* males. AVT is upregulated in dominant *A. burtoni* males (Greenwood et al., 2008; Renn et al., 2008) and dominant individuals of both sexes in *N. pulcher* (Aubin-Horth et al., 2007), a relative of *Julidochromis*. Furthermore, manipulation of AVT alters territorial behavior in the tropical damselfish (Santangelo and Bass, 2006). These studies establish a clear relationship between AVT and aggressive behavior, and support the hypothesis that AVT plays an important role in regulating aggressive behavior in *J. marlieri*, the most aggressive phenotype in our study.

RT-qPCR verification

We selected genes for verification of array results using qPCR on both RNA and amplified RNA from the *Julidochromis* individuals. There was good correlation between results from RT-qPCR and microarray data for parvalbumin ($F = 4.611$, $P = 0.05732$) and isotocin ($F = 51.3$, $P = 3.06 \times 10^{-5}$). However, RT-qPCR results for MAP-1B

differed from microarray results substantially. The correlation found between the amplified RNA and total RNA RT-qPCR results for isotocin ($F = 6.728$, $P = 0.0268$) and parvalbumin ($F = 4.511$, $P = 0.05962$) shows that the amplification process does not significantly change the detection of relative expression. Given the lack of correlation between amplified RNA and total RNA for MAP-1B, in addition to the lack of correlation with microarray results, we suspect that either the primers or array features or both are showing non-specific effects.

Meta-analysis for novel function inference

Perhaps the most exciting future direction for comparative microarray studies is the potential to investigate the identified gene modules in light of other studies and to use these results to infer biological function and identify novel candidates. Nearly half of the genes found in the sex-specific or phenotype-specific modules have no sequence information, while others have no known homolog in GenBank (Table 3). Array studies such as this one provide a tentative functional annotation for many of the genes of unknown function that are on the array (Landry and Aubin-Horth, 2007). The current study identified a number of new genes that are likely to be important in aggressive behavior and parental care in *Julidochromis* and other cichlids, which previously had not been identified (see Table S1 in supplementary material). To see whether any of the putative dominance-related genes in this study had been previously identified as aggression-related genes, we compared our data with a previous study conducted with a smaller microarray comparing *A. burtoni* territorial males, non-territorial males and females (Renn et al., 2008). We predicted that because the polygynous *A. burtoni* has conventional sex-specific behavior, its pattern of sex-specific

Table 3. Number of significantly upregulated genes in each phenotype with sequence information and those with no known homolog in GenBank

Group	Number of genes upregulated ($P < 0.05$)	Number of genes with sequence information	Number of genes without a known homolog
<i>J. transcriptus</i> males	412	196	50
<i>J. transcriptus</i> females	301	136	63
<i>J. marlieri</i> females	529	216	85
<i>J. marlieri</i> males	356	179	41
Dominance related	76	34	5
Subordinance related	24	16	6
Female related	24	10	3
Male related	21	10	2

gene expression may align more closely with that of the conventional *J. transcriptus* rather than the reversed *J. marlieri*. Unexpectedly, we found that *J. marlieri* females showed a trend towards sharing significant gene regulation with *A. burtoni* males ($P=0.0966$), while no other phenotype had significantly greater shared regulation with dominant *A. burtoni* males than expected by random chance. In addition to AVT, 14 genes with no functional information were upregulated in these two aggressive phenotypes. In addition to gonadotropin releasing hormone (GnRH), prolactin and somatotropin, male *J. transcriptus* expressed four genes with no functional information that were also upregulated in dominant *A. burtoni*. Two of these were part of the dominance module also being upregulated in *J. marlieri* females. Genes that are associated with multiple aggressive phenotypes are likely to be important in aggressive behavior and will be especially exciting to pursue in future investigations. These genes may have key regulatory roles in the behavior of cichlids and other organisms but, because they have not been previously characterized, they have not been formally investigated. The results of this analysis reiterate how many genes of ecological importance are unknown, and how microarray studies can contribute to providing tentative annotations for these genes (Landry and Aubin-Horth, 2007).

It is interesting to note that *J. marlieri* females share more gene expression with *A. burtoni* territorial males than the male aggressive phenotype in *J. transcriptus*. This latter fact leads to little overlap between the *Julidochromis* dominance module and genes upregulated in territorial *A. burtoni* males. One interesting possibility to explain the patterns of overlap between these modules involves the temporal control of territoriality in *A. burtoni*. Male *A. burtoni* cycle between territorial and non-territorial states, requiring behavioral plasticity in aggression and dominance. Significant overlap between *A. burtoni* males and *J. marlieri* females may suggest that aggression in *J. marlieri* females was initially a plastic response, more similar to that seen in *A. burtoni* males, that has become fixed over evolutionary time. In contrast, ancestral male dominance as seen in *J. transcriptus* may be controlled by different mechanisms.

Evolutionary implications of gene-expression patterns

The phylogenetic relationship of species in the genus *Julidochromis* has been the subject of recent debate. While *Julidochromis* was previously considered to be a monophyletic group, recent phylogenetic hypotheses, based on both mitochondrial and nuclear sequence data, suggest paraphyly (Sturmbauer et al., 2010) as well as hybridization leading to the preservation of ancestral alleles. While the exact

evolutionary relationship of *Julidochromis* species remains unresolved the most recent phylogeny suggests that *J. transcriptus* and *J. marlieri* are ~3.7 million years diverged (Sturmbauer et al., 2010) but it remains clear that female aggression in *J. marlieri* is the derived characteristic, raising the issue of sex-role reversal. Aggressive and territorial behaviors in biparental species function to defend valuable territories, protect young from predators and prevent extra-pair copulations. Male territory defense in most cichlids may be adaptive due to the fact that sexual size dimorphism is typically male biased (Erlandsson and Ribbinck, 1997). Increased female size in *J. marlieri* could have evolved as a result of increased sexual selection on females (Blanckenhorn, 2005) and female aggression may have been a facultative response. Regardless of the particular driving forces, it is clear that *J. marlieri* has undergone significant changes in sex-specific behavior since *J. marlieri* and *J. transcriptus* diverged. Our results, which show limited conserved sex-specific gene regulation between *J. transcriptus* and *J. marlieri*, support the conclusion that major changes in sex-specific gene expression have accompanied this behavioral divergence.

Despite divergence in sex-specific gene expression, there was significant overlap in phenotype-related genes between species, particularly those associated with aggressive behavior. Our identification of a core set of genes shared among aggressive phenotypes fits with the recent evidence leading to the theory of a 'genetic toolkit' for social behavior. This theory, similar to ideas concerning the evolution of development, suggests that there may be a small set of conserved genes specialized or co-opted to produce certain behaviors (Toth et al., 2007). Complex behaviors, such as social behavior, could theoretically be constructed from altering this underlying set of behaviorally specialized genes. Evidence for this concept has been found in comparisons of a similar but independently evolved foraging behavior in honeybees and paper wasps that are ~100 million years diverged (Toth et al., 2010). Researchers found that relatively few genes showed phenotype-specific expression across species but suggested that these genes may be major regulators of behavior. Although *J. transcriptus* and *J. marlieri* are more closely related than the taxa discussed above, the core set of genes strongly associated with aggressive behavior in *J. transcriptus* and *J. marlieri* conforms to the genetic toolkit theory. The fact that the aggressive phenotype in *J. marlieri* females is independently evolved reinforces the concept of a genetic toolkit for aggressive behavior, and presents a candidate group of genes important in regulating aggression in Tanganyikan cichlids. Although comparisons with *A. burtoni* reveal that this genetic toolkit may not include all Tanganyikan cichlids, such comparisons are complicated by aggressive plasticity in *A. burtoni*. Further comparative studies and additional time points will be needed in order to determine whether the identified core set of aggression genes is in fact used consistently to regulate aggressive behavior in a range of species.

Another theme common in developmental biology that can influence the study of behavior is the concept that changes in traits can often result from changes not only in genes but in gene expression (Carroll, 2000; Toth et al., 2007) (but see Hoekstra and Coyne, 2007). Given that there are species-level differences between *J. transcriptus* and *J. marlieri* in aggressive behavior it will be interesting to directly compare levels of aggression-related genes between *Julidochromis* species.

CONCLUSIONS

While *J. transcriptus* follows the ancestral behavioral pattern of male-biased dominance and female-biased parental care behavior,

J. marlieri is reversed in these sex-specific behaviors. We predicted that there would be a high degree of overlap in sex-specific and phenotype-specific gene expression between *J. marlieri* and *J. transcriptus* due to their relatedness and similarities in phenotype-specific behavior. However, our investigation revealed that there has been significant divergence in gene expression patterns between *J. marlieri* and *J. transcriptus*, perhaps due to the reversal of sex-biased behavior in *J. marlieri*. The degree to which these differences orchestrate sex-specific behavior as opposed to other divergent phenotypic traits needs to be addressed in future functional work. While most comparisons revealed small overlap between species, overlap between the dominant phenotypes was larger and highly significant. Closer examination of gene expression patterns (Fig. 4) shows that a substantial proportion of the most highly regulated genes in *J. transcriptus* males and *J. marlieri* females are part of the dominance module. The large overlap in dominance genes between *J. transcriptus* and *J. marlieri* suggests that the gene expression patterns underlying aggressive behavior are shared between the two species, raising the possibility of a genetic toolkit for aggressive behavior in Tanganyikan cichlids. In addition, we found that a number of candidate genes previously associated with aggressive behavior were strongly associated with the aggressive phenotypes in each species. We further provided a tentative functional annotation for a number of unknown genes that are associated with aggressive phenotypes. Investigating the roles of the unknown genes in these phenotype-specific modules is a crucial next step in understanding the mechanisms of aggressive, subordinate and sex-specific behavior in Tanganyikan cichlids.

LIST OF ABBREVIATIONS

AVT	arginine vasotocin
GABA	γ -Aminobutyric acid
GAPDH	glyceraldehyde 3-phosphate dehydrogenase
GEL50	gene expression level 50
GHRH	growth hormone-releasing hormone
GnRH	gonadotrophin-releasing hormone
GSI	gonadosomatic index
LIMMA	Linear Models for Microarray normalization data
MAP-1B	microtubule-associated protein-1B
RT-qPCR	real-time quantitative PCR

ACKNOWLEDGEMENTS

We thank Victoria Zero, Kelsey Wood and Heather Machado for insightful discussion on the design and execution of the project. They, and members of the Aubin-Horth lab, provided comment on earlier drafts of this manuscript. This work was funded by NIH: R15 GM080727-01A1 to S.C.P.R. as well as a grant from the James F. and Marion L. Miller Foundation to M.E.S. Deposited in PMC for release after 12 months.

REFERENCES

- Aubin-Horth, N., Desjardins, J. K., Martei, Y. M., Balshine, S. and Hofmann, H. A. (2007). Masculinized dominant females in a cooperatively breeding species. *Mol. Ecol.* **16**, 1349-1358.
- Awata, S., Takeuchi, H. and Kohda, M. (2006). The effect of body size on mating system and parental roles in a biparental cichlid fish (*Julidochromis transcriptus*): a preliminary laboratory experiment. *J. Ethol.* **24**, 125-132.
- Barlow, G. W. and Lee, J. S. F. (2005). Sex-reversed dominance and aggression in the cichlid fish *Julidochromis marlieri*. *Ann. Zool. Fenn.* **42**, 477-483.
- Bender, N., Taborsky, M. and Power, D. H. (2008). The role of prolactin in the regulation of brood care in the cooperatively breeding fish *Neolamprologus pulcher*. *J. Exp. Zool.* **309A**, 515-524.
- Berglund, A. and Rosenqvist, G. (2003). Sex-role reversal in pipefish. *Adv. Stud. Behav.* **32**, 131-167.
- Blackenhorn, W. U. (2005). Behavioral causes and consequences of sexual size dimorphism. *Ethology* **111**, 977-1016.
- Blumstein, D. T. and Daniel, J. C. (2007). *Quantifying Behavior the JWatcher Way*. Sunderland, MA: Sinauer Associates, Inc. www.jwatcher.ucla.edu.
- Bridges, R. S., Numan, M., Ronsheim, P. M., Mann, P. E. and Lupini, C. E. (1990). Central prolactin infusions stimulate maternal behavior in steroid treated, nulliparous female rats. *Proc. Natl. Acad. Sci. USA* **87**, 8003-8007.

- Buitenhuis, B., Hedegaard, J., Janss, L. and Sorensen, P. (2009). Differentially expressed genes for aggressive pecking behaviour in laying hens. *BMC Genomics* **10**, 544.
- Burger, J. (1981). Sexual differences in parental activities of breeding Black Skimmers. *Am. Nat.* **117**, 6.
- Burmeister, S. S. and Fernald, R. D. (2005). Evolutionary conservation of the egr-1 immediate-early gene response in a teleost. *J. Comp. Neurol.* **481**, 220-232.
- Carroll, S. B. (2000). Endless forms: the evolution of gene regulation and morphological diversity. *Cell* **101**, 577-580.
- Cello, M. R. (1986). Parvalbumin in most gamma-aminobutyric acid-containing neurons of the rat cerebral cortex. *Science* **231**, 995-997.
- Churchill, G. A. (2002). Fundamentals of experimental design for cDNA microarrays. *Nat. Genet.* **32**, 490-495.
- Clark, T. A. and Townsend, J. P. (2007). Quantifying variation in gene expression. *Mol. Ecol.* **16**, 2613-2616.
- Clutton-Brock, T. H. and Vincent, A. C. J. (1991). Sexual selection and the potential reproductive rates of males and females. *Nature* **351**, 58-60.
- Cohen, R., Chalifa-Caspi, V., Williams, T. D., Auslander, M., George, S. G., Chipman, J. K. and Tom, M. (2007). Estimating the efficiency of fish cross-species cDNA microarray hybridization. *Mar. Biotechnol.* **9**, 491-499.
- Creelman, E. and Storey, E. H. (1991). Sex differences in reproductive behavior of Atlantic Puffins. *Condor* **93**, 390-398.
- Cummings, M. E., Larkins-Ford, J., Reilly, C. R. L., Wong, R. Y., Ramsey, M. and Hoffman, H. A. (2008). Sexual and social stimuli elicit rapid and contrasting genomic responses. *Proc. R. Soc. Lond. B* **275**, 393-402.
- de Almeida, R. M. M., Ferrari, P. H., Parmigiani, S. and Miczek, K. A. (2005). Escalated aggressive behavior: dopamine, serotonin and GABA. *Eur. J. Pharmacol.* **526**, 51-64.
- Derome, N. and Bernatchez, L. (2006). The transcriptomics of ecological convergence between two limnetic coregonine fishes (Salmonidae). *Mol. Biol. Evol.* **23**, 2370-2378.
- Dusenbery, D. B. (1980). Chemotactic behavior of mutants of the nematode *Caenorhabditis elegans* that are defective in osmotic avoidance. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* **137**, 93-96.
- Edwards, A. C., Rollman, S. M., Morgan, T. J. and Mackay, T. F. C. (2006). Quantitative genomics of aggressive behavior in *Drosophila melanogaster*. *PLoS Genet.* **2**, e154.
- Eens, M. and Pinxten, R. (2000). Sex-role reversal in vertebrates: behavioural and endocrinological accounts. *Behav. Processes* **51**, 135-147.
- Ellegren, H. and Parsch, J. (2007). The evolution of sex-biased genes and sex-biased gene expression. *Nat. Rev. Genet.* **8**, 689-698.
- Emlen, S. T. and Wrege, P. H. (2004). Size dimorphism, intrasexual competition, and sexual selection in wattled jacana (*Jacana jacana*), a sex-role-reversed shorebird in Panama. *Auk* **121**, 391-403.
- Erlandsson, A. and Ribbink, A. J. (1997). Patterns of sexual size dimorphism in African cichlid fishes. *S. Afr. J. Sci.* **93**, 498-508.
- Filby, A. L., Paull, G. C., Hickmoew, T. F. A. and Tyler, C. R. (2010). Unravelling the neurophysiological basis of aggression in a fish model. *BMC Genomics* **11**, 498-514.
- Foran, C. M. and Bass, A. H. (1999). Preoptic GnRH and AVT: axes for sexual plasticity in teleost fish. *Gen. Comp. Endocrinol.* **116**, 141-152.
- Forsgren, E., Amundsen, T., Borg, A. and Bjelvenmark, J. (2004). Unusually dynamic sex roles in a fish. *Nature* **429**, 551-554.
- Fraser, G. S., Jones, I. L. and Hunter, F. M. (2002). Male-female differences in parental care in monogamous Crested Auklets. *Condor* **104**, 413-423.
- Gammie, S. C., Auger, A. P., Jessen, H. M., Vanzo, R. J., Awad, T. A. and Stevenson, S. A. (2007). Altered gene expression in mice selected for high maternal aggression. *Genes Brain Behav.* **6**, 432-443.
- Garcia, V., Jouventin, P. and Mauget, R. (1996). Parental care and the Prolactin secretion pattern in the king penguin: an endogenously timed mechanism? *Horm. Behav.* **30**, 259-265.
- Goodwin, N. B., Balshine-Earn, S. and Reynolds, J. D. (1998). Evolutionary transitions in parental care in cichlid fish. *Proc. R. Soc. Lond. B* **265**, 2265-2272.
- Greenwood, A. K., Wark, A. R., Fernald, R. D. and Hofmann, H. A. (2008). Expression of arginine vasotocin in distinct preoptic regions is associated with dominant and subordinate behavior in an African cichlid fish. *Proc. Biol. Sci.* **275**, 2392-2402.
- Gwynne, D. T. (1985). Role-reversal in katydids: habitat influences reproductive behaviour (Orthoptera: Tettigoniidae, *Metaballus* sp.). *Behav. Ecol. Sociobiol.* **16**, 355-361.
- Hoekstra, H. E. and Coyne, J. A. (2007). The locus of evolution: Evo Devo and the genetics of adaptation. *Evolution* **61**, 995-1016.
- Insel, T. R. and Young, L. J. (2000). Neuropeptides and the evolution of social behavior. *Curr. Opin. Neurobiol.* **10**, 784-789.
- Itzkowitz, M. (1984). Parental division of labor in a monogamous fish. *Behaviour* **89**, 251-260.
- Itzkowitz, M., Santangelo, N. and Richter, M. (2001). Parental division of labour and the shift from minimal to maximal role specializations: an examination using a biparental fish. *Anim. Behav.* **61**, 1237-1245.
- Itzkowitz, M., Santangelo, N. and Richter, M. (2003). How does a parent respond when its mate emphasizes the wrong role? A test using monogamous fish. *Anim. Behav.* **66**, 863-869.
- Itzkowitz, M., Santangelo, N., Cleveland, A., Bockelman, A. and Richter, M. (2005). Is the selection of sex-typical parental roles based on an assessment process? A test in the monogamous convict cichlid fish. *Anim. Behav.* **69**, 95-105.
- Jin, W., Riley, R. M., Wolfinger, R. D., White, K. P., Passador-Gurel, G. and Gibson, G. (2001). The contributions of sex, genotype and age to transcriptional variance in *Drosophila melanogaster*. *Nat. Genet.* **29**, 389-395.
- Jones, A. G., Walker, D. and Avise, J. C. (2001). Genetic evidence for extreme polyandry and extraordinary sex-role reversal in a pipefish. *Proc. R. Soc. Lond. B* **268**, 2531-2535.

- Keenleyside, M. H. A. and Bletz, B. F. (1981). The reproductive behavior of *Aequidens vittatus* (Pisces, cichlidae) in Surinam, South America. *Environ. Biol. Fishes* **6**, 87-94.
- Konings, A. (1998). *Tanganyikan Cichlids in their Natural Habitat*, pp. 111-126. Cichlid Press. ISBN 0-9668255-0-0.
- Konopka, R. J. and Benzer, S. (1971). Clock mutants of *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **68**, 2112-2116.
- Kuwamura, T. (1986). Parental care and mating systems of cichlid fishes in Lake Tanganyika: a preliminary field study. *J. Ethol.* **4**, 129-146.
- Kuwamura, T., Kawanabe, H., Hori, M. and Nagoshi, M. (1997). The evolution of parental care and mating systems among Tanganyikan cichlids. In *Fish Communities in Lake Tanganyika* (ed. H. Kawanabe, M. Hori and M. Nagoshi), pp. 57-86. Kyoto: Kyoto University Press.
- Landry, C. R. and Aubin-Horth, N. (2007). Ecological annotation of genes and genomes through ecological genomics. *Mol. Ecol.* **16**, 4419-4421.
- Lavery, R. J. and Reeb, S. J. (1994). Effect of mate removal on current and subsequent parental care in the convict cichlid (Pisces: Cichlidae). *Ethology* **97**, 265-277.
- Lewohl, J. M., Wang, L., Miles, M. P., Zhang, L. H., Dodd, P. R. and Harris, A. (2000). Gene expression in human alcoholism: microarray analysis of frontal cortex. *Alcohol. Clin. Exp. Res.* **24**, 1873-1882.
- Machado, H. E., Pollen, A. A., Hoffman, H. A. and Renn, S. C. P. (2009). Interspecific profiling of gene expression informed by comparative genomic hybridization: a review and a novel approach in African cichlid fishes. *Integr. Comp. Biol.* **49**, 644-659.
- Marivoet, S., Moons, L. and Vandesande, F. (1988). Localization of growth hormone releasing factor-like immunoreactivity in the hypothalamo-hypophyseal system of the frog (*Rana temporaria*) and the sea bass (*Dicentrarchus labrax*). *Gen. Comp. Endocrinol.* **72**, 72-79.
- Mayer, I., Rosenqvist, G., Borg, B., Anhenjo, I., Berlung, A. and Schulz, R. W. (1993). Plasma levels of sex steroids in three species of pipefish (Syngnathidae). *Can. J. Zool.* **71**, 1903-1907.
- Millesi, E., Hoffman, I. E., Steurer, S., Metwaly, M. and Dittami, J. P. (2002). Vernal changes in the behavior and endocrine responses to GnRH application in male European ground squirrels. *Horm. Behav.* **41**, 51-58.
- Osborne, K. A., Robichon, A., Burgess, E., Butland, S., Shaw, R. A., Coulthard, A., Pereira, H. S., Greenspan, R. J. and Sokolowski, M. B. (1997). Natural behavior polymorphism due to a cGMP-dependent protein kinase of *Drosophila*. *Science* **277**, 834-836.
- Price, E. O., Adams, T. E., Huxsoll, C. C. and Borgwardt, R. E. (2003). Aggressive behavior is reduced in bulls actively immunized against gonadotropin-releasing hormone. *J. Anim. Sci.* **81**, 411-415.
- R Development Core Team (2006). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Renn, S. C. P., Aubin-Horth, N. and Hofmann, H. A. (2004). Biologically meaningful expression profiling across species using heterologous hybridization to a cDNA microarray. *BMC Genomics* **5**, 42.
- Renn, S. C. P., Aubin-Horth, N. and Hofmann, H. A. (2008). Fish and chips: functional genomics of social plasticity in an African cichlid fish. *J. Exp. Biol.* **211**, 3041-3056.
- Rhodes, J. S. and Crabbe, J. C. (2005). Gene expression induced by drugs of abuse. *Curr. Opin. Pharmacol.* **5**, 26-33.
- Robinson, G. E., Fernald, R. D. and Clayton, D. F. (2008). Genes and social behavior. *Science* **322**, 896-900.
- Salzburger, W., Renn, S. C. P., Steinke, D., Braasch, I., Hoffman, H. A. and Meyer, A. (2008). Annotation of expressed sequence tags for the East African cichlid fish *Astatotilapia burtoni* and evolutionary analysis of cichlid ORFs. *BMC Genomics* **9**, 96-110.
- Santangelo, N. and Bass, A. H. (2006). New insights into neuropeptide modulation of aggression: field studies of arginine vasotocin in a territorial tropical damselfish. *Proc. R. Soc. Lond. B* **273**, 3085-3092.
- Santos, E. M., Kille, P., Workman, V., Paull, G. C. and Tyler, C. R. (2008). Sexually dimorphic gene expression in the brains of mature zebrafish. *Comp. Biochem. Physiol.* **149**, 314-324.
- Schradin, C. and Anzenberger, G. (1999). Prolactin, the hormone of paternity. *News Physiol. Sci.* **14**, 223-231.
- Schumer, M. (2009). Gene expression, hormones, and behavior in a sex-role conventional and sex-role reversed cichlid species pair (*Julidochromis*). BA thesis, Reed College, Portland, OR, USA.
- Sen Sarma, M., Rodriguez-Zas, S. L., Hong, F., Zhong, S. and Robinson, G. E. (2009). Transcriptomic profiling of the central nervous system regions in three species of the honey bee during dance communication behavior. *PLoS ONE* **4**, e6408.
- Smyth, G. K. (2004). Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3**, Article 3.
- Soma, K. K., Francis, R. C., Wingfield, J. C. and Fernald, R. D. (1996). Androgen regulation of hypothalamic neurons containing gonadotropin-releasing hormone in a cichlid fish: integration with social cues. *Horm. Behav.* **30**, 216-226.
- Sturmhuber, C., Salzburger, W., Duffer, N., Schelly, R. and Koblmüller, S. (2010). Evolutionary history of the Lake Tanganyikan cichlid tribe Lamprologini (Teleostei: Perciformes) derived from mitochondrial and nuclear DNA data. *Mol. Phylogenet. Evol.* **57**, 266-284.
- Toth, A. L. and Robinson, G. E. (2007). Evo-devo and the evolution of social behavior. *Trends Genet.* **23**, 334-341.
- Toth, A. L., Varala, K., Henshaw, M. T., Rodriguez-Zas, S. L., Hudson, M. E. and Robinson, G. E. (2010). Brain transcriptomic analysis in paper wasps identifies genes associated with behavior across social insect lineages. *Proc. R. Soc. Lond. B* **277**, 2139-2148.
- Trivers, R. (1972). Parental investment and sexual selection. In *Sexual Selection and the Descent of Man 1871-1971* (ed. B. Campbell), pp. 136-207. Chicago: Aldine.
- Voigt, C. and Goymann, W. (2008). Sex-role reversal is reflected in the brain of African black coucals (*Centropus grillii*). *Dev. Neurobiol.* **67**, 1560-1573.
- White, S. A. and Fernald, R. D. (1993). Gonadotropin-releasing hormone-containing neurons change size with reproductive state in female *Haplochromis burtoni*. *J. Neurosci.* **13**, 434-441.
- Whitfield, C. W., Cziko, A. M. and Robinson, G. E. (2003). Gene expression profiles in the brain predict behavior in individual honey bees. *Science* **302**, 296-299.
- Wisenden, B. D., Lanfranconi-Izawa, T. L. and Keenleyside, M. H. A. (1995). Fin digging and leaf lifting by the convict cichlid, *Cichlasoma nigrofasciatum*: examples of parental food provisioning. *Anim. Behav.* **49**, 623-631.
- Yamagishi, S. and Kohda, M. (1996). Is the cichlid fish *Julidochromis marlieri* polyandrous? *Ichthyol. Res.* **43**, 469-471.
- Yamamoto, D., Jallon, J. M. and Komatsu, A. (1997). Genetic dissection of sexual behavior in *Drosophila melanogaster*. *Annu. Rev. Entomol.* **42**, 551-585.
- Yang, X., Schadt, E. E., Wang, S., Arnold, A. P., Ingram-Drake, L., Drake, T. A. and Lusis, A. J. (2006). Tissue-specific expression and regulation of sexually dimorphic genes in mice. *Genome Res.* **16**, 995-1004.