

Section 9 Regression with Stationary Time Series

How time-series regression differs from cross-section

- Natural ordering of observations contains information
 - Random reshuffling of observations would obscure dynamic economic relationship, but leave traditional regression unchanged
 - How can we incorporate this dynamic information into our regression model?
- We usually think of the data as being drawn from a potentially infinite **data-generating process** rather than from a finite population of observations.
- Variables are often call “time series” or just “series” rather than variables
 - Index observations by time period t
 - Number of observations = T
- Dynamic relationship means that not all of the effects of x_t occur in period t .
 - A change in x_t is likely to affect y_{t+1} , y_{t+2} , etc.
 - By the same logic, y_t depends not only on x_t but also on x_{t-1} , x_{t-2} , etc.
 - We model these dynamic relationships with **distributed lag** models, in which
$$y_t = f(x_t, x_{t-1}, x_{t-2}, \dots).$$
- We will need to focus on the dynamic elements of both the deterministic relationship between the variables and the stochastic relationship (error term)
- The dynamic ordering of observations means that the error terms are usually **serially correlated** (or autocorrelated over time)
 - Shocks to the regression are unlikely to completely disappear before the following period
 - Exception: stock market returns, where investors should respond to any shock and make sure that next period’s return is not predictable
 - Two observations are likely to be more highly correlated if they are close to the same time than if they are more widely separated.
 - Covariance matrix of error term will have non-zero off-diagonal elements, with elements lying closest to the diagonal likely being substantially positive and decreasing as one moves away from the diagonal.
- **Nonstationary** time series create problems for econometrics.
 - We will study implications of and methods for dealing with nonstationarity in Section 12.
 - Example will illustrate nature of problem (“spurious regressions”)
 - Regression of AL attendance on Botswana real GDP
 - Correlation = 0.9656

- $R^2 = 0.9323$
- Coefficient has t of 24.90.
- Good regression?

Source	SS	df	MS	Number of obs = 47		
Model	3.4342e+15	1	3.4342e+15	F(1, 45)	=	619.89
Residual	2.4930e+14	45	5.5400e+12	Prob > F	=	0.0000
				R-squared	=	0.9323
				Adj R-squared	=	0.9308
Total	3.6835e+15	46	8.0077e+13	Root MSE	=	2.4e+06

ALAttend	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
rgdpl2	3285.11	131.9447	24.90	0.000	3019.36	3550.86
_cons	8029710	640681.3	12.53	0.000	6739311	9320108

- Correlation is spurious because both series are trending upward, so most of each series' deviation from mean is due to separate trends.
- Much of the last 20 years in econometrics has been devoted to understanding how to deal with nonstationary time series.
- We will study this intensively in a few weeks.
- Nonstationarity forces us to remove the common trend (often by differencing) before interpreting the correlation or regression

Lag operators and differences

- With time-series data we are often interested in the relationship among variables at different points in time.
- Let x_t be the observation corresponding to time period t .
 - The first lag of x is the preceding observation: x_{t-1} .
 - We sometimes use the **lag operator** $L(x_t)$ or $Lx_t \equiv x_{t-1}$ to represent lags.
 - We often use higher-order lags: $L^s x \equiv x_{t-s}$.
- The first difference of x is the difference between x and its lag:
 - $\Delta x_t \equiv x_t - x_{t-1} = (1 - L)x_t$
 - Higher-order differences are also used:

$$\Delta^2 x_t = \Delta(\Delta x_t) = (x_t - x_{t-1}) - (x_{t-1} - x_{t-2}) = x_t - 2x_{t-1} + x_{t-2}$$

$$= (1 - L)^2 x_t = (1 - 2L + L^2)x_t$$
 - $\Delta^s x_t = (1 - L)^s x_t$
- Difference of the log of a variable is approximately equal to the variable's growth rate:

$$\Delta(\ln x_t) = \ln x_t - \ln x_{t-1} = \ln(x_t / x_{t-1}) \approx x_t / x_{t-1} - 1 = \Delta x_t / x_t$$
 - Log difference is exactly the continuously-compounded growth rate
 - The discrete growth-rate formula $\Delta x_t / x_t$ is the formula for once-per-period compounded growth

- Lags and differences in Stata
 - First you must define the data to be time series: `tsset year`
 - This will correctly deal with missing years in the year variable.
 - Can define a variable for quarterly or monthly data and set format to print out appropriately.
 - For example, suppose your data have a variable called `month` and one called `year`. You want to combine into a single time variable called `time`.
 - `gen time = ym(year, month)`
 - This variable will have a `%tm` format and will print out like `2010m4` for April 2010.
 - You can then do `tsset time`
 - Once you have the time variable set, you can create lags with the lag operator `L` and differences with `D`.
 - For example, last period's value of `x` is `L.x`
 - The change in `x` between now and last period is `D.x`
 - Higher-order lags and differences can be obtained with `L3.x` for third lag or `D2.x` for second difference.

Autocovariance and autocorrelation

- Autocovariance of order s is $\text{cov}(x_t, x_{t-s})$
 - We generally assume that the autocovariance depends only on s , not on t .
 - This is analogous to our Assumption #0: that all observations follow the same model (or were generated by the same data-generating process)
 - This is *one element* of a time series being stationary
- Autocorrelation of order s (ρ_s) is the correlation coefficient between x_t and x_{t-s} .
 - $$\rho_k = \frac{\text{cov}(x_t, x_{t-k})}{\text{var}(x_t)}$$
 - We estimate with
$$r_k = \frac{\frac{1}{T-k} \sum_{t=k+1}^T (x_t - \bar{x})(x_{t-k} - \bar{x})}{\frac{1}{T} \sum_{t=1}^T (x_t - \bar{x})^2}$$
 - We sometimes subtract one from both denominators, or sometimes ignore the different fractions in front of the summations since their ratio goes to 1 as T goes to ∞ .
- ρ_k as a function of k is called the **autocorrelation function** of the series and its plot is often called a **correlogram**.

Some simple univariate time-series models

- We sometimes represent a variable's time-series behavior with a univariate model.

- **White noise:** The simplest univariate time-series process is called white noise $y_t = v_t$, where v_t is a mean-zero IID error (usually normal).
 - The key point here is the autocorrelations of white noise are all zero (except, of course, for ρ_0 , which is always 1).
 - Very few economic time series are white noise.
 - Changes in stock prices are probably one.
 - We use white noise as a basic building block for more useful time series:
 - Consider problem of forecasting y_t conditional on all past values of y .
 - $y_t = E[y_t | y_{t-1}, y_{t-2}, \dots] + v_t$
 - Since any part of the past behavior of y that would help to predict the current y should be accounted for in the expectation part, the error term v should be white noise.
 - The one-period-ahead forecast error of y should be white noise.
 - We sometimes call this forecast-error series the “fundamental underlying white noise series for y ” or the “innovations” in y .
- The simplest autocorrelated series is the **first-order autoregressive (AR(1)) process**: $y_t = \beta_0 + \beta_1 y_{t-1} + v_t$, where e is white noise.
 - In this case, our one-period-ahead forecast is $E[y_t | y_{t-1}] = \beta_0 + \beta_1 y_{t-1}$ and the forecast error is v_t .
 - For simplicity, suppose that we have removed the mean from y so that $\beta_0 = 0$.
 - Consider the effect of a one-time shock v_1 on the series y from time one on, assuming (for simplicity) that $y_0 = 0$ and all subsequent v values are also zero.
 - $y_1 = \beta_1(0) + v_1 = v_1$
 - $y_2 = \beta_1 y_1 + v_2 = \beta_1 v_1$
 - $y_3 = \beta_1 y_2 + v_3 = \beta_1^2 v_1$
 - $y_s = \beta_1^{s-1} v_1$.
 - This shows that the effect of the shock on y “goes away” over time only if $|\beta_1| < 1$.
 - The condition $|\beta_1| < 1$ is necessary for the AR(1) process to be **stationary**.
 - If $\beta_1 = 1$, then shocks to y are permanent. This series is called a **random walk**.
 - The random walk process can be written $y_t = y_{t-1} + v_t$ or $\Delta y_t = v_t$. The first difference of a random walk is stationary and is white noise.

- If y follows a stationary AR(1) process, then $\rho_1 = \beta_1, \rho_2 = \beta_1^2, \dots, \rho_s = \beta_1^s$.
 - One way to attempt to identify the appropriate specification for a time-series variable is to examine the autocorrelation function of the series.
 - If the autocorrelation function declines exponentially toward zero, then the series might follow an AR(1) process with positive β_1 .
 - A series with $\beta_1 < 0$ would oscillate back and forth between positive and negative responses to a shock.
 - The autocorrelations would also oscillate between positive and negative while converging to zero.

Assumptions of time-series regression

- Before we deal with issues of specifications of y and x , we will think about the problems that serially correlated error terms cause for OLS regression. (GHL's Section 9.3)
- Can estimate time-series regressions by OLS as long as y and x are stationary and x is exogenous.
 - **Exogeneity:** $E(e_t | x_t, x_{t-1}, \dots) = 0$.
 - **Strict exogeneity:** $E(e_t | \dots, x_{t+2}, x_{t+1}, x_t, x_{t-1}, x_{t-2}, \dots) = 0$.
- Assumptions of time-series regression:
 - **TSMR2:** y and x are stationary and x is strictly exogenous
 - **TSMR3:** $E(e_t) = 0$
 - **TSMR4:** $\text{var}(e_t) = \sigma^2$
 - **TSMR5:** $\text{cov}(e_t, e_s) = 0, t \neq s$
 - **TSMR6:** $e_t \sim N(0, \sigma^2)$
- However, nearly all time-series regressions are prone to having serially correlated error terms, which violates TSMR5.
 - Omitted variables are probably serially correlated
- This is a particular form of violation of the IID assumption.
 - Observations are correlated with those of nearby periods
- As long as the other OLS assumptions are satisfied, this causes a problem not unlike heteroskedasticity
 - OLS is still unbiased and consistent
 - OLS is not efficient
 - OLS estimators of standard errors are biased, so cannot use ordinary t statistics for inference
- To some extent, adding more lags of y and x to the specification can reduce the severity of serial correlation.

- Two methods of dealing with serial correlation of the error term:
 - GLS regression in which we transform the model to one whose error term is not serially correlated
 - This is analogous to weighted least squares (also a GLS procedure)
 - Estimate by OLS but use standard error estimates that are robust to serial correlation

Detecting autocorrelation

- We can test the autocorrelations of a series to see if they are zero.
 - Asymptotically, $\sqrt{T}r_k \sim N(\rho_k, 1)$, so we can compute this as a test statistic and test against the null hypothesis $\rho_k = 0$.
- **Breusch-Godfrey Lagrange multiplier test** for autocorrelation:
 - Regress y (or residuals) on x and lagged residuals (first-order, or more)
 - Use F test of residual coefficient(s) in y regression or TR^2 in residual regression as χ^2
- **Box-Ljung Q test** for null hypothesis that the first k autocorrelations are zero:

$$Q_k = T(T+2) \sum_{j=1}^k \frac{r_j^2}{T-j}$$
 is asymptotically χ_k^2 .
- **Durbin-Watson test** used to be the standard test for first-order autocorrelation, but was difficult because critical values depend on x . Not used much anymore.

Estimation with autocorrelated errors

- OLS with autocorrelated errors
 - Assumption TSMR4 is violated, which leads to inefficient estimators and biased standard errors just like in case of heteroskedasticity
 - **Important special case:** We will see that a common distributed lag model puts y_{t-1} on the right-hand side as a regressor. This causes special problems when there is serial correlation because
 - e_{t-1} is part of y_{t-1}
 - e_{t-1} is correlated with e_t
 - Therefore e_t is correlated with one of the regressors, which leads to bias and inconsistency in the coefficient estimators.
 - If we can transform the model into one that has no autocorrelation (for example, v_t if error term is $e_t = \rho e_{t-1} + v_t$), then we can get consistent OLS estimators as long as all the x variables are exogenous (but not necessarily strictly exogenous) with respect to v .
- **HAC consistent standard errors** (Newey-West)

- As with White's heteroskedasticity consistent standard errors, we can correct the OLS standard errors for autocorrelation as well.
- We know that

$$b_2 = \beta_2 + \frac{\frac{1}{T} \sum_{i=1}^T (x_i - \bar{x}) e_i}{\frac{1}{T} \sum_{i=1}^T (x_i - \bar{x})^2}.$$

- In this formula, $\text{plim } \bar{x} = \mu_X$, $\text{plim} \left(\frac{1}{T} \sum_{i=1}^T (x_i - \bar{x})^2 \right) = \sigma_X^2$.

- So $\text{plim}(b_2 - \beta_2) = \frac{\text{plim} \left(\frac{1}{T} \sum_{i=1}^T (x_i - \mu_X) e_i \right)}{\sigma_X^2} = \frac{\text{plim}(\bar{u})}{\sigma_X^2}$, where $\bar{u} = \frac{1}{T} \sum_{i=1}^T u_i$ and

$$u_i \equiv (X_i - \mu_X) e_i.$$

- And in large samples, $\text{var}(b_2) = \text{var} \left(\frac{\bar{u}}{\sigma_X^2} \right) = \frac{\text{var}(\bar{u})}{\sigma_X^4}$.

- Under IID assumption, $\text{var}(\bar{u}) = \frac{1}{T} \text{var}(u_i) = \frac{\sigma_u^2}{T}$, and the formula reduces to one we know from before.
- However, serial correlation means that the error terms are not IID (and x is usually not either), so this doesn't apply.

- In the case where there is serial correlation we have to take into account the covariance of the u_i terms:

$$\begin{aligned} \text{var}(\bar{u}) &= \text{var} \left(\frac{u_1 + u_2 + \dots + u_T}{T} \right) \\ &= \frac{1}{T^2} \left[\sum_{i=1}^T \sum_{j=1}^T E(u_i u_j) \right] \\ &= \frac{1}{T^2} \sum_{i=1}^T \left(\text{var}(u_i) + \sum_{j \neq i} \text{cov}(u_i, u_j) \right) \\ &= \frac{1}{T^2} \left[T \text{var}(u_i) + 2(T-1) \text{cov}(u_i, u_{i-1}) + 2(T-2) \text{cov}(u_i, u_{i-2}) + \dots + 2 \text{cov}(u_i, u_{i-(T-1)}) \right] \\ &= \frac{\sigma_u^2}{T} f_T, \end{aligned}$$

where

$$\begin{aligned} f_T &\equiv 1 + 2 \sum_{j=1}^{T-1} \left(\frac{T-j}{T} \right) \text{corr}(u_i, u_{i-j}) \\ &= 1 + 2 \sum_{j=1}^{T-1} \left(\frac{T-j}{T} \right) \rho_j. \end{aligned}$$

- Thus, $\text{var}(b_2) = \left[\frac{1}{T} \frac{\sigma_u^2}{\sigma_x^4} \right] f_T$, which expresses the variance as the product of the no-autocorrelation variance and the f_T factor that corrects for autocorrelation.
- In order to implement this, we need to know f_T , which depends on the autocorrelations of u for orders 1 through $T - 1$.
 - These are not known and must be estimated.
 - For ρ_1 we have lots of information because there are $T - 1$ pairs of values for (u_t, u_{t-1}) in the sample.
 - For ρ_{T-1} , there is only one pair $(u_t, u_{t-(T-1)})$ —namely (u_T, u_1) —on which to base an estimate.
 - The **Newey-West** procedure truncates the summation in f_T at some value $m - 1$, so we estimate the first $m - 1$ autocorrelations of v using the OLS residuals and compute $\hat{f}_T = 1 + 2 \sum_{j=1}^{m-1} \left(\frac{m-j}{m} \right) r_j$.
 - m must be large enough to provide a reasonable correction but small enough relative to T to allow the r values to be estimated well.
 - Stock and Watson suggest choosing $m = 0.75T^{\frac{1}{3}}$ as a reasonable rule of thumb.
- To implement in Stata, use hac option in xtreg (with panel data) or post-estimation command newey , lags(m)
- **GLS with an AR(1) error term**
 - One of the oldest time-series models (and not used so much anymore) is the model in which e_t follows an AR(1) process:

$$y_t = \beta_0 + \beta_1 x_t + e_t$$

$$e_t = \rho e_{t-1} + v_t,$$
 where v is a white-noise error term and $-1 < \rho < 1$.
 - In practice, $\rho > 0$ nearly always
 - GLS transforms the model into one with an error term that is not serially correlated.
 - Let

$$\tilde{y}_t = \begin{cases} y_t \sqrt{(1-\rho^2)}, & t=1, \\ y_t - \rho y_{t-1}, & t=2, 3, \dots, T, \end{cases} \quad \tilde{x}_t = \begin{cases} x_t \sqrt{(1-\rho^2)}, & t=1, \\ x_t - \rho x_{t-1}, & t=2, 3, \dots, T, \end{cases}$$

$$\tilde{e}_t = \begin{cases} e_t \sqrt{(1-\rho^2)}, & t=1, \\ e_t - \rho e_{t-1}, & t=2, 3, \dots, T. \end{cases}$$
 - Then $\tilde{y}_t = (1-\rho)\beta_1 + \beta_2 \tilde{x}_t + \tilde{e}_t$.

- The error term in this regression is equal to v_t for observations 2 through T and is a multiple of e_1 for the first observation.
- By assumption, v is white noise and values of v in periods after 1 are uncorrelated with e_1 , so there is no serial correlation in this transformed model.
- If the other assumptions are satisfied, it can be estimated efficiently by OLS.
- But what is ρ ?
 - Need to estimate ρ to calculate feasible GLS estimator.
 - Traditional estimator for ρ is $\hat{\rho} = \text{corr}(\hat{e}_t, \hat{e}_{t-1})$ using OLS residuals.
 - This estimation can be iterated to get a new estimate of ρ based on the GLS estimator and then re-do the transformation: repeat until converged.
- Two-step estimator using FGLS based on $\hat{\rho}$ is called the **Prais-Winsten** estimator (or **Cochrane-Orcutt** when first observation is dropped).
- Problems: ρ is not estimated consistently if e_t is correlated with x_t , which will always be the case if there is a lagged dependent variable present and may be the case if x is not strongly exogenous.
 - In this case, we can use nonlinear methods to estimate ρ and β jointly by search.
 - This is called the **Hildreth-Lu** method.
- In Stata, the prais command implements all of these methods (depending on option). Option corc does Cochrane-Orcutt; ssearch does Hildreth-Lu; and the default is Prais-Winsten.
- You can also estimate this model with ρ as the coefficient on y_{t-1} in an OLS model, with or without the restriction implied in HGL's equation (9.44).

Distributed-lag models

- Modeling the **deterministic part** of a dynamic relationship between two variables
- In general, the distributed-lag model has the form $y_t = \alpha + \sum_{i=0}^{\infty} \beta_i x_{t-i} + e_t$. But of course, we cannot estimate an infinite number of lag coefficients β_i , so we must either truncate or find another way to approximate an infinite lag structure.
 - **Multipliers**
 - $\frac{\partial E(y_t)}{\partial x_{t-s}} = \beta_s$ is the s -period delay multiplier, telling how much a one-time, temporary shock to x would be affecting y s periods later
 - β_0 is the “impact multiplier”

- $\sum_{r=0}^s \beta_r$ is the cumulative or “interim” multiplier after s periods. It measures the cumulative effect of a permanent change in x on y s periods later.
- $\sum_{s=1}^{\infty} \beta_s$ is the “total multiplier” measuring the final cumulative effect of a permanent change in x on y .
- We can easily have additional regressors with either the same or different lag structures.
- **Finite distributed lags**
 - $y_t = \alpha + \beta_0 x_t + \beta_1 x_{t-1} + \dots + \beta_q x_{t-q} + e_t = \alpha + \beta(L)x_t + e_t$
 - This is finite distributed-lag model of order q
 - Under assumptions TSMR1–6, the model can be estimated by OLS. If there is serial correlation, then the appropriate correction must be made to standard errors or a GLS model must be used.
 - Problems with finite DL model
 - If x is strongly autocorrelated, then collinearity will be a problem and it will be difficult to estimate individual lag weights accurately
 - Difficult to know appropriate lag length (can use AIC or SC)
 - If lags are long and sample is short, will lose lots of observations.
- **Koyck lag: y is AR(1) with regressors**
 - $y_t = \delta + \theta_1 y_{t-1} + \delta_0 x_t + v_t$
 - $\frac{\partial y_t}{\partial x_t} = \delta_0$
 - $\frac{\partial y_{t+1}}{\partial x_t} = \theta_1 \delta_0$
 - $\frac{\partial y_{t+2}}{\partial x_t} = \theta_1^2 \delta_0$
 - $\frac{\partial y_{t+s}}{\partial x_t} = \theta_1^s \delta_0$

Thus, dynamic multipliers start at δ_0 and decay exponentially to zero over infinite time. Thus, this is effectively a distributed lag of infinite length, but with only 2 parameters (plus intercept) to estimate.

 - Cumulative multipliers are $\sum_{k=0}^s \frac{\partial y_{t+k}}{\partial x_t} = \delta_0 \sum_{k=0}^s \theta_1^k$.
 - Long-run effect of a permanent change is $\delta_0 \sum_{k=0}^{\infty} \theta_1^k = \frac{\delta_0}{1 - \theta_1}$.
 - Estimation has the potential problem of **inconsistency** if v_t is serially correlated.

- This is a serious problem, especially as some of the test statistics for serial correlation of the error are biased when the lagged dependent variable is present.
 - Koyck lag is parsimonious and fits lots of lagged relationships well.
 - With multiple regressors, the Koyck lag applies the same lag structure (rate of decay) to all regressors.
 - Is this reasonable for your application?
 - Example: delayed adjustment of factor inputs: can't stop using expensive factor more quickly than you start using cheaper factor.
- **ARX(p) Model**
 - We can generalize the Koyck lag model to longer lags:

$$y_t = \delta + \theta_1 y_{t-1} + \dots + \theta_p y_{t-p} + \delta_0 x_t + v_t.$$
 - This can be written $\theta(L)y_t = \delta + \delta_0 x_t + v_t$.
 - Same general principles apply:
 - Worry about stationarity of lag structure: roots of $\theta(L)$
 - If v is serially correlated, OLS will be biased and inconsistent
 - Dynamic multipliers are determined by coefficients of infinite lag polynomial $[\theta(L)]^{-1}$
 - If more than on x , all have same lag structure
 - How to determine length of lag p ?
 - Can keep adding lags as long as θ_p is statistically significant
 - Can choose to max the Akaike information criterion (AIC) or Bayesian (Schwarz) information criterion (SC).
 - Note that regression can use as many as $T - p$ observations, but should use the same number for all regressions with different p values in assessing information criteria.
 - AIC will choose longer lag than SC.
 - AIC came first, so is still used a lot
 - SC is asymptotically unbiased
 - Stata calculates info criteria by estat ic (after regression)
- **ADL(p, q) Model: "Rational" lag**
 - We can also add lags to the x variable(s)
 - $y_t = \delta_0 + \theta_1 y_{t-1} + \dots + \theta_p y_{t-p} + \delta_0 x_t + \delta_1 x_{t-1} + \dots + \delta_q x_{t-q} + v_t$
 - Can add more x variables with varying lag lengths
 - $\theta(L)y_t = \delta_0 + \delta(L)x_t + v_t,$
 - $y_t = \frac{\delta_0}{\theta(L)} + \frac{\delta(L)}{\theta(L)}x_t + \frac{v_t}{\theta(L)}.$

- Multipliers are the (infinite) coefficients on the lag polynomial $\frac{\delta(L)}{\theta(L)}$
- Stationarity depends only on $\theta(L)$, not on $\delta(L)$.
- Can easily estimate this by OLS assuming:
 - $E(v_t | y_{t-1}, y_{t-2}, \dots, y_{t-p}, x_t, x_{t-1}, \dots, x_{t-q}) = 0$
 - (y_t, x_t) has same mean, variance, and autocorrelations for all t
 - (y_t, x_t) and (y_{t-s}, x_{t-s}) become independent as $s \rightarrow \infty$
 - No perfect multicollinearity
- These are general TSMR assumptions that apply to most time-series models.

Forecasting with time-series models

- **With AR(p) model**
 - $y_t = \delta + \theta_1 y_{t-1} + \theta_2 y_{t-2} + \dots + \theta_p y_{t-p} + v_t$, with v assumed to be serially uncorrelated.
 - We have observations for $t = 1, 2, \dots, T$
 - Forecast for $T+1$: $\hat{y}_{T+1} = \hat{\delta} + \hat{\theta}_1 y_T + \hat{\theta}_2 y_{T-1} + \dots + \hat{\theta}_p y_{T-p+1}$ with expected value of v at zero because of no serial correlation
 - If error term were autoregressive, then conditional expectation of v_{T+1} would be ρv_T , so would include residual of observation T
 - Forecast error is $u_1 = y_{T+1} - \hat{y}_{T+1} = (\delta - \hat{\delta}) + \sum_{s=1}^p (\theta_s - \hat{\theta}_s) y_{T+1-s} + v_{T+1}$
 - HGL assume that $\text{var}(v_{T+1}) \gg \text{var}\left[(\delta - \hat{\delta}) + \sum_{s=1}^p (\theta_s - \hat{\theta}_s) y_{T+1-s}\right]$, so they ignore the latter.
 - I'm not willing to ignore this (although it is often true that the variance of the error is larger, as in Problem 6.17.
 - I will write $e_{b,k} = (\delta - \hat{\delta}) + \sum_{s=1}^p (\theta_s - \hat{\theta}_s) y_{T+k-s}$ and $\text{var}(e_{b,k})$ to be that component of the k -period-ahead forecast and keep it in the equation
 - $u_1 = e_{b,1} + v_{T+1}$,
 - $\text{var}(u_1) = \text{var}(e_{b,1}) + \sigma_v^2$
 - What about forecast for $T+2$?
 - y_{T+1} appears on the right-hand side of the $T+2$ equation, so we substitute our one-period-ahead forecast of it:

$$\hat{y}_{T+2} = \hat{\delta} + \hat{\theta}_1 \hat{y}_{T+1} + \hat{\theta}_2 y_T + \dots + \hat{\theta}_p y_{T+2-p}$$

- Forecast error is

$$\begin{aligned} u_2 &= y_{T+2} - \hat{y}_{T+2} = (\delta - \hat{\delta}) + \theta_1 (\hat{y}_{T+1} - y_{T+1}) + \sum_{s=1}^p (\theta_s - \hat{\theta}_s) y_{T+2-s} + v_{T+2} \\ &= e_{b,2} + \theta_1 e_{b,1} + v_{T+2} + \theta_1 v_{T+1}. \end{aligned}$$

- Variance of forecast error is

$$\text{var}(u_2) = \text{var}(e_{b,2}) + \theta_1^2 \text{var}(e_{b,1}) + (1 + \theta_1^2) \sigma_v^2.$$

- Similarly,

$$\text{var}(u_3) = \text{var}(e_{b,3}) + \theta_1^2 \text{var}(e_{b,2}) + (\theta_1^2 + \theta_2^2) \text{var}(e_{b,1}) + (1 + \theta_1^2 + (\theta_2 + \theta_1^2)^2) \sigma_v^2$$