

Ali Pardini, Jonah Rohlfing, and Danielle Cass





#### Metal Uptake in Slough Sedge





- Car brake wear and use is characterized by heavy metal pollution; specifically Cu and Zn emissions (1)
- In freshwater systems, elevated levels of Cu and Zn in the cells of fish damages certain enzymes responsible for mediating oxidative stress (2)
- Plants in bioswales, like slough sedge, are capable of accumulating heavy metals from roads into their tissues, preventing metals from entering water systems (3)

How effective are the slough sedges of the Reed College bioswale on 28th st. at accumulating Cu and Zn from their surroundings?

### Methods for Sample Collections and Digestion


#### Plant sample collection





From nine sections of the bioswale, 45 samples of the roots, above ground plant material and soil were taken.

# Soil sample collection map



#### Digesting soil, root, and above ground plant material samples with HNO<sub>3</sub> and H<sub>2</sub>O<sub>2</sub>



Samples heated at 75°C until dry and ground to a fine powder



Samples digested for 12 hours with 500 μL of 70% metal free HNO<sub>3</sub>



500 μL of H<sub>2</sub>O<sub>2</sub> added to samples and set on 50°C hot plate



Samples placed in the microwave digester for 10 minutes at 180°C



Samples filtered into a falcon tube and diluted to 10 mL with 1% HNO<sub>3</sub>




Now the samples are ready for the ICP-MS!


### Methods for Metals Analysis: AAS or ICP-MS

#### **Atomic Absorption Spectra**

AAS detects elements through the absorption of characteristic wavelengths of light (4).




Zn standard curve, detection limit < 50 ppb



## **Inductively Couples Plasma Mass Spectrometry**

The ICP-MS uses plasma to ionize the elements in a sample and then measures the ions using a mass spectrometer (5).



	(mg/kg)	(mg/kg)	(%)
Rye grass Cu	10.2	9.03	12.95
Rye grass Zn	30.5	18.3	6.69
Buffalo soil Zn	408	120.9	237.2

Expected Experimental % error

#### Conclusions

- Analysis of NIST standards shows that a more effective soil digestion method is required
- The ICP-MS is the preferred instrument for analysis of Cu and Zn

#### **Moving Forward**

- Using ICP-MS to quantity the levels of Cu and Zn in each sample
- Determine the effectiveness of the bioswale

#### Acknowledgments

Many thanks to the staff and faculty of the Reed College Chemistry Department.
Particular thanks to Danielle Cass and Randy Hicks for their help and leadership.

#### **Literature Cited**

review. Environ Sci Pollut Res Int. 2015 Feb;22(4):2491-504. doi: 10.1007/s11356-014-3696-8. E 2014 Oct 17. PMID: 25318420; PMCID: PMC4315878.

Latif, F., Iqbal, R., Ambreen, F., Kousar, S., Ahmed, T., & Aziz, S. (2024). Studies on bioaccumulation patterns, biochemical and genotoxic effects of copper on freshwate fish, catla catla: An in vivo analysis. [Estudos sobre padro de bioacumulação, efeitos bioquímicos e genotóxicos do cobre em peixes de água doce, Catla catla: Uma análise vivo] Brazilian Journal of Biology, 84 doi:10.1590/1519-6984.256905

- Wyatt, G., and S. J. Matiasek. "Seasonal Variation in F Bioaccumulation of Heavy Metals at the Butte College Bioswale." NASA/ADS, https://ui.adsabs.harvard.edu/a 2019AGUFM.H41J1838W/abstract.
- "Atomic Absorption Spectrometry (AAS) Information."
   Thermo Fisher Scientific US, https://
   www.thermofisher.com/us/en/home/industrial/spectrosc
   elemental-isotope-analysis/spectroscopy-elemental-isotope-analysis/spectroscopy-elemental-isotope-analysis-learning-center/trace-elemental-analysis-tea-information/atomic-absorption-aa-information.html#:~:text=Atomic%20absorption%20spemetry%20(AAS)
   %20detects,absorbances%20are%20measured%20ag
- "A Beginner's Guide to ICP-MS, ICP-MS Analysis and E Mass Spectrometry." Chemical Analysis, Life Sciences, Diagnostics, https://www.agilent.com/en/support/atomic spectroscopy/inductively-coupled-plasma-massspectrometry-icp-ms/icp-ms-instruments/what-is-icp-ms