

**MATHEMATICS 211**

ASSIGNMENT 5

Due: October 8, 2014

01° Let  $A$  be an antisymmetric matrix:

$$A = \begin{pmatrix} 0 & -w & v \\ w & 0 & -u \\ -v & u & 0 \end{pmatrix}$$

Compute the determinant of  $A$ .

02° Let  $a$  be a real number, distinct from 0. Let  $f$  be the mapping carrying  $\mathbf{R}^3 \setminus \{0\}$  to  $\mathbf{R} = \mathbf{R}^1$ , defined as follows:

$$f(x, y, z) := (x^2 + y^2 + z^2)^a$$

Find the value(s) of  $a$  for which:

$$f_{xx}(x, y, z) + f_{yy}(x, y, z) + f_{zz}(x, y, z) = 0$$

03° Consider the following curve in  $\mathbf{R}^3$ :

$$\Gamma(t) := (\exp(t)\cos(t), \exp(t)\sin(t), t)$$

where  $t$  is any real number (soit *time*). Find the angle between the position vector  $\Gamma(t)$  and the velocity vector  $\Gamma'(t)$  at time  $t = \pi/4$ .

04° Let  $f$  be the real valued function defined on  $\mathbf{R}^3$  as follows:

$$f(x, y, z) = z - (x^2 + y^2) \quad ((x, y, z) \in \mathbf{R}^3)$$

Let  $M$  be the level set in  $\mathbf{R}^3$  defined by the relation:

$$f(x, y, z) = 0$$

Clearly, the (position) vector  $(1, 2, 5)$  lies in  $M$ . The tangent plane  $T_{(1,2,5)}(M)$  to  $M$  at  $(1, 2, 5)$  consists of the vectors  $(u, v, w)$  in  $\mathbf{R}^3$  which meet the condition:

$$(f_x(1, 2, 5), f_y(1, 2, 5), f_z(1, 2, 5)) \bullet (u, v, w) = d$$

where  $d$  is a suitable number. Find  $d$ .

05° Let  $F$  be the mapping carrying  $\mathbf{R}^2$  to  $\mathbf{R}^3$  defined by the following relations:

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = F\left(\begin{pmatrix} u \\ v \end{pmatrix}\right) : \begin{array}{l} x = u \\ y = v \\ z = u^2 + v^2 \end{array}$$

Let  $M$  be the range of  $F$ . Describe the tangent plane:

$$T_P(M)$$

to  $M$  at the point:

$$P = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} = F\left(\begin{pmatrix} 1 \\ 2 \end{pmatrix}\right)$$

By definition, the vectors in  $T_P(M)$  have the form:

$$\begin{pmatrix} p \\ q \\ r \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \begin{pmatrix} \circ & \bullet \\ \circ & \bullet \\ \circ & \bullet \end{pmatrix} \begin{pmatrix} u-1 \\ v-2 \end{pmatrix}$$

where:

$$\begin{pmatrix} \circ & \bullet \\ \circ & \bullet \\ \circ & \bullet \end{pmatrix} = DF\left(\begin{pmatrix} 1 \\ 2 \end{pmatrix}\right)$$

and where:

$$\begin{pmatrix} u \\ v \end{pmatrix}$$

runs through all vectors in  $\mathbf{R}^2$ . Find a vector:

$$N = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$$

which is perpendicular to  $T_P(M)$ . In fact, you can take  $N$  to be:


$$\begin{pmatrix} \circ \\ \circ \\ \circ \end{pmatrix} \times \begin{pmatrix} \bullet \\ \bullet \\ \bullet \end{pmatrix}$$

Draw a diagram to illustrate the sense of this problem.

06° Let  $H$  be the Hipparchus Map:

$$H\left(\begin{pmatrix} \phi \\ \theta \end{pmatrix}\right) = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \cos\theta\cos\phi \\ \cos\theta\sin\phi \\ \sin\theta \end{pmatrix}$$

where  $\phi$  is the longitude and  $\theta$  is the latitude. In the following picture of the range  $\mathbf{S}^2$  of  $H$ :



plot the vector:

$$V = H \begin{pmatrix} \pi/4 \\ \pi/6 \end{pmatrix} + DH \begin{pmatrix} \pi/4 \\ \pi/6 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Be ye exact.

07° Let  $a$  and  $b$  be any numbers for which  $0 < b < a$ . Let  $c$  be the positive number which satisfies the relation:  $b^2 + c^2 = a^2$ . Let  $f$  be the function defined on  $\mathbf{R}^2$  as follows:

$$f(x, y) = \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2$$

Let  $(p, q)$  be any member of  $\mathbf{R}^2$  for which  $f(p, q) = 1$ . [The set of all such members  $(p, q)$  compose an *ellipse* in  $\mathbf{R}^2$ . We will make a drawing of it in the lectures.] Let  $\alpha$  be the angle between the vectors:

$$(p, q) - (-c, 0) \quad \text{and} \quad (f_x(p, q), f_y(p, q))$$

and let  $\beta$  be the angle between the vectors:

$$(p, q) - (+c, 0) \quad \text{and} \quad (f_x(p, q), f_y(p, q))$$

Show that  $\alpha = \beta$ . This result explains the phenomenon of the Whispering Gallery.

08° Find the equation of the tangent plane for the surface:

$$S : \quad \sqrt{x} + \sqrt{y} + \sqrt{z} = 4 \quad (0 < x, 0 < y, 0 < z)$$

at the point  $(1, 4, 1)$ .