

MATHEMATICS 211
ASSIGNMENT 11
Due: December 3, 2014

01• Memorize the Greek alphabet:

α	alpha	A
β	beta	B
γ	gamma	Γ
δ	delta	Δ
ϵ	epsilon	E
ζ	zeta	Z
η	eta	H
θ	theta	Θ
ι	iota	I
κ	kappa	K
λ	lambda	Λ
μ	mu	M
ν	nu	N
ξ	xi	Ξ
o	omicron	O
π	pi	Π
ρ	rho	P
σ	sigma	Σ
τ	tau	T
v	upsilon	Υ
ϕ	phi	Φ
χ	chi	X
ψ	psi	Ψ
ω	omega	Ω

02° Let ϕ be a scalar field and let F be a vector field on \mathbf{R}^3 . By definition, ϕ is a function for which the domain is (a suitable subset of) \mathbf{R}^3 and the codomain is \mathbf{R} :

$$\phi(x, y, z)$$

while F is a function for which the domain is (a suitable subset of) \mathbf{R}^3 and the codomain is \mathbf{R}^3 :

$$F(x, y, z) = (u(x, y, z), v(x, y, z), w(x, y, z))$$

where u , v , and w are (in effect) scalar fields, the components of F .

One defines the following operators acting on ϕ and F :

(•) **Gradient**

$$\nabla\phi = (\partial\phi/\partial x, \partial\phi/\partial y, \partial\phi/\partial z)$$

(•) **Curl**

$$\nabla \times F = (\partial w/\partial y - \partial v/\partial z, \partial u/\partial z - \partial w/\partial x, \partial v/\partial x - \partial u/\partial y)$$

(•) **Divergence**

$$\nabla \bullet F = \partial u/\partial x + \partial v/\partial y + \partial w/\partial z$$

(•) **Laplacian**

$$\nabla^2\phi = \nabla \bullet (\nabla\phi) = \partial^2\phi/\partial x^2 + \partial^2\phi/\partial y^2 + \partial^2\phi/\partial z^2$$

Show that:

$$\nabla \times (\nabla\phi) = (0, 0, 0) \quad \text{and} \quad \nabla \bullet (\nabla \times F) = 0$$

03° Given vector fields G and H on \mathbf{R}^3 , show that:

$$\nabla \bullet (G \times H) = (\nabla \times G) \bullet H - G \bullet (\nabla \times H)$$

04° Given a vector field F on \mathbf{R}^3 , show that:

$$\nabla \times (\nabla \times F) = \nabla(\nabla \bullet F) - \nabla^2 F$$

Of course, ∇^2 acts on F component by component.

05° Let ϕ be the scalar field on \mathbf{R} defined as follows:

$$\phi(x, y, z) = -\frac{1}{r} \quad (0 < r)$$

where:

$$r = \sqrt{x^2 + y^2 + z^2}$$

Calculate:

$$-(\nabla\phi)(x, y, z)$$

To that end, note that $\partial r/\partial x = x/r$, $\partial r/\partial y = y/r$, and $\partial r/\partial z = z/r$.

06° Let D be a subset of \mathbf{R}^3 and let F be a vector field on \mathbf{R}^3 defined on D :

$$F(x, y, z) = (u(x, y, z), v(x, y, z), w(x, y, z)) \quad ((x, y, z) \in D)$$

Let J be a closed finite interval in \mathbf{R} :

$$J = [a, b] \quad (a < b)$$

and let Γ be a (parametrized) curve in \mathbf{R}^3 defined on J :

$$\Gamma(t) = (x(t), y(t), z(t)) \quad (a \leq t \leq b)$$

Let the range of Γ be included in the domain of F :

$$\Gamma(t) \in D \quad (a \leq t \leq b)$$

In this context, one defines the *line integral* of F over Γ as follows:

$$\begin{aligned} \int_{\Gamma} F &:= \int_a^b F(\Gamma(t)) \bullet \Gamma^\circ(t) dt \\ &= \int_a^b [(u(x(t), y(t), z(t)))x^\circ(t) + (v(x(t), y(t), z(t)))y^\circ(t) + (w(x(t), y(t), z(t)))z^\circ(t)] dt \end{aligned}$$

For the following particular cases:

$$F(x, y, z) = \left(-\frac{y}{x^2 + y^2}, \frac{x}{x^2 + y^2}, z\right) \quad (0 < x^2 + y^2)$$

and:

$$\Gamma_1(t) = (\cos(t), \sin(t), t), \quad \Gamma_2(t) = (\cos(t), -\sin(t), t) \quad (0 \leq t \leq \pi)$$

show that:

$$(\nabla \times F)(x, y, z) = (0, 0, 0)$$

and:

$$\int_{\Gamma_1} F \neq \int_{\Gamma_2} F$$

Note, however, that the initial points of Γ_1 and Γ_2 coincide. The same is true of the terminal points. Finally, with reference to the preceding problem, replace F by:

$$F(x, y, z) = -(\nabla\phi)(x, y, z)$$

Show that, in this case:

$$\int_{\Gamma_1} F = \int_{\Gamma_2} F$$

07° Let α be a function defined on the interval $\mathbf{R}^+ := (0, \infty)$ in \mathbf{R} and let ϕ be the scalar field defined on the region $D := \mathbf{R}^3 \setminus \{(0, 0, 0)\}$ in \mathbf{R}^3 , as follows:

$$\phi(x, y, z) = \alpha(r) \quad (0 < r := \sqrt{x^2 + y^2 + z^2})$$

Show that:

$$(\nabla\phi)(x, y, z) = \alpha'(r) \frac{1}{r}(x, y, z) \quad (0 < r := \sqrt{x^2 + y^2 + z^2})$$

08° Let β be a function defined on the interval $\mathbf{R}^+ := (0, \infty)$ in \mathbf{R} and let F be the vector field defined on the region $D := \mathbf{R}^3 \setminus \{(0, 0, 0)\}$ in \mathbf{R}^3 , as follows:

$$F(x, y, z) = \beta(r)(x, y, z) \quad (0 < r := \sqrt{x^2 + y^2 + z^2})$$

Show that:

$$(\nabla \times F)(x, y, z) = (0, 0, 0) \quad (0 < r := \sqrt{x^2 + y^2 + z^2})$$

Find a scalar field ϕ for which:

$$F(x, y, z) = (\nabla\phi)(x, y, z) \quad (0 < r := \sqrt{x^2 + y^2 + z^2})$$

To do so, apply the foregoing problem. Work out the details for the following case:

$$\beta(r) = r^a \quad (0 < r)$$

where a is any real number.

09° Let J be an interval in \mathbf{R} and let D be a region in \mathbf{R}^3 . Let ϕ be a scalar field “defined on D but depending on t ”:

$$\phi(t, x, y, z) \quad (t \in J, (x, y, z) \in D)$$

One defines the following operator acting on ϕ :

(•) **d’Alembertian**

$$\square^2\phi = \partial^2\phi/\partial t^2 - \nabla^2\phi = \partial^2\phi/\partial t^2 - \partial^2\phi/\partial x^2 - \partial^2\phi/\partial y^2 - \partial^2\phi/\partial z^2$$

Let G and H be vector fields “defined on D but depending on t ”:

$$G(t, x, y, z), \quad H(t, x, y, z) \quad (t \in J, (x, y, z) \in D)$$

and satisfying the following relations:

$$\begin{aligned} \nabla \bullet G &= 0, & \nabla \bullet H &= 0 \\ \partial G/\partial t - \nabla \times H &= (0, 0, 0), & \partial H/\partial t + \nabla \times G &= (0, 0, 0) \end{aligned}$$

Show that:

$$\square^2 G = (0, 0, 0) \quad \text{and} \quad \square^2 H = (0, 0, 0)$$

Of course, $\partial/\partial t$ and \square^2 act on G and H component by component. Conclude that any one of the components of G and H , let it be ϕ , satisfies the **wave equation**:

$$\square^2 \phi = 0$$